

## Product datasheet for **MR207249L3V**

### Vrk2 (BC013520) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Vrk2 (BC013520) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Vrk2
Synonyms:	2810003O05Rik; A1447698
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	BC013520
ORF Size:	1362 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR207249).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">BC013520</a> , <a href="#">AAH13520</a>
RefSeq Size:	1473 bp
RefSeq ORF:	1364 bp
Locus ID:	69922
Cytogenetics:	11 A3.3



[View online »](#)

**Gene Summary:**

Serine/threonine kinase that regulates several signal transduction pathways. Isoform 1 modulates the stress response to hypoxia and cytokines, such as interleukin-1 beta (IL1B) and this is dependent on its interaction with MAPK8IP1, which assembles mitogen-activated protein kinase (MAPK) complexes. Inhibition of signal transmission mediated by the assembly of MAPK8IP1-MAPK complexes reduces JNK phosphorylation and JUN-dependent transcription. Phosphorylates histone H3. Phosphorylates 'Thr-18' of p53/TP53, and thereby increases its stability and activity. Phosphorylates BANF1 and disrupts its ability to bind DNA and reduces its binding to LEM domain-containing proteins. Downregulates the transactivation of transcription induced by ERBB2, HRAS, BRAF, and MEK1. Blocks the phosphorylation of ERK in response to ERBB2 and HRAS. May also phosphorylate MAPK8IP1. Can also phosphorylate the following substrates that are commonly used to establish in vitro kinase activity: casein, MBP and histone H2B, but it is not sure that this is physiologically relevant (By similarity).[UniProtKB/Swiss-Prot Function]