

## Product datasheet for **MR207405**

### Igh (BC031470) Mouse Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Igh (BC031470) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Igh
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin
ORF Nucleotide Sequence:	>MR207405 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGACTTGAGACTGAGCTGTGCTTTTATTATTGTTCTTTTAAAAGGGTCCAGAGTGAAGTGAAGCTTG  
AGGAGTCTGGAGGAGCTTGGTGAACCTGGAGGATCCATGAAACTCTCCTGTAGCCTCTGGATTAC  
TTTCAGTAACTCCTGGATGAAGTGGTCCGCCAGTCTCCAGAGAAGGGCTTGAGTGGTGTCTCAAATT  
AGATTGAGATCTGATAATTATGCAACACATTATGCGGAGTCTGTGAAAGGGAGGTTACCATCTCAAGAG  
ATGATTCCAAAAGTCGTCTCTACCTGCAATGAGCAGTTAAGGGCTGAAGACTGGAATTTACTG  
CACAAATGCTATGGACTACTGGGTCAAGGAACCTCAGTCACCGTCTCCTCAGCCAAAACAACAGCCCCA  
TCGGTCTATCCACTGGCCCTGTGTGGAGATACAACCTGGTCTCGGTGACTTAGGATGCCTGGTCA  
AGGGTATTTCCCTGAGCCAGTGACCTTGACCTGGAACCTGGATCCCTGTCCAGTGGTGTGCACACCTT  
CCCAGTGTCTGCAGTCTGACCTCTACACCCTCAGCAGCTCAGTACTGTAACCTCGAGCACCTGGCCC  
AGCCAGTCCATCACCTGCAATGTGGCCACCCGGCAAGCAGCACCAAGGTGGACAAGAAAATTGAGCCCA  
GAGGGCCCAATCAAGCCCTGTCTCCATGCAAATGCCAGCACCTAACCTCTTGGGTGGACCATCCGT  
CTTCATCTTCCCTCCAAAGATCAAGGATGACTCATGATCTCCCTGAGCCCCATGGTCACATGTGTGGT  
GTGGATGTGAGCGAGGATGACCCAGATGTCAGATCAGCTGGTTCGTGAACAACGTGGAAGTACTCACAG  
CTCAGACACAAACCCATAGAGAGGATTACAACAGTACTCTCCGGGTGGTCAAGTGCCTCCCCATCCAGCA  
CCAGGACTGGATGAGTGGCAAGGAGTTCAAATGCAAGGTCAACAACAAAGCCCTCCCAGCGCCCATCGAG  
AGAACCATCTCAAACCCAAAGGGTCAAGAGCTCCACAGGTATATGTCTTGCCTCCACCAGAAGAAG  
AGATGACTAAGAAACAGGTCACCTGACCTGCATGGTACAGACTTATGCCTGAAGACATTTACGTGGA  
GTGGACCAACAACGGGAAAACAGAGCTAACTACAAGAACACTGAACAGTCTGGACTCTGATGGTTCT  
TACTTCATGTACAGCAAGCTGAGAGTGGAAAAGAAGAACTGGGTGGAAAGAAATAGTACTCTGCTCAG  
TGGTCCACGAGGGTCTGCACAATCACACACACTAAGAGCTTCTCCCGACTCCGGGTAAA

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)


**Protein Sequence:** >MR207405 protein sequence  
 Red=Cloning site Green=Tags(s)

MDLRLSCAFIIVLLKGVQSEVKLEESGGGLVQPGGSMKLSCVASGFTFSNSWMNWVRQSPEKGLEWVAQI  
 RLRSDNYATHYAESVKGRFTISRDDSKSRLYLQMSLRAEDTGIYYCTNAMYWGQGTSVTVSSAKTTAP  
 SVYPLAPVCGD TTGSSVTLGCLVKGYFPEPVTLTWNSGSLSSGVHTFPAVLQSDLYTLSSSVTVTSSTWP  
 SQSITCNVAHPASSTKVDK KIEPRGPTIKPCPPCKPAPNLLGGPSVFI FPPKIKDVLMI LSPMVT CVV  
 VD VSEDDPDVQISW FVNNVEVLTAQTQTHREDYNSTLRVVSALPIQH QDWM S GKEFKCKVNNKALPAPIE  
 RTISKPKG SVRAPQVYVLP PPEEEMTKQVTLTCMVTDFMPEDIYVEWTNNGKTELNYKNTEPVLDS DGS  
 YFMYSKLRVEKKNWVERNSYSCSVVHEGLHNHHTTKSFSRTPGK


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**Plasmid Map:**



<b>ACCN:</b>	BC031470
<b>ORF Size:</b>	1392 bp
<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">BC031470</a>
<b>RefSeq Size:</b>	1658 bp
<b>RefSeq ORF:</b>	1394 bp
<b>Locus ID:</b>	111507
<b>Cytogenetics:</b>	12 F1-F2 12
<b>MW:</b>	51.2 kDa

**Gene Summary:**

Summary: Immunoglobulins recognize foreign antigens and initiate immune responses such as phagocytosis and the complement system. Each immunoglobulin molecule consists of two identical heavy chains and two identical light chains. This region represents the germline organization of the heavy chain locus. The locus includes V (variable), D (diversity), J (joining), and C (constant) segments. During B cell development, a recombination event at the DNA level joins a single D segment with a J segment; this partially rearranged D-J gene is then joined to a V segment. The rearranged V-D-J is then transcribed with the IGHM constant region; this transcript encodes a mu heavy chain. Later in development B cells generate V-D-J-Cmu-Cdelta pre-messenger RNA, which is alternatively spliced to encode either a mu or a delta heavy chain. Mature B cells in the lymph nodes undergo switch recombination, so that the V-D-J gene is brought in proximity to one of the IGHG, IGHA, or IGHE genes and each cell expresses either the gamma, alpha, or epsilon heavy chain. Recombination of many different V segments with several J segments provides a wide range of antigen recognition. Additional diversity is attained by junctional diversity, resulting from the random additional of nucleotides by terminal deoxynucleotidyltransferase, and by somatic hypermutation, which occurs during B cell maturation in the spleen and lymph nodes. The RefSeq represents the IGH locus from C57BL/6. Several V and D segments in C57BL/6 are known to be incapable of encoding a protein and are considered pseudogenes. [provided by RefSeq, Jul 2008]