

Product datasheet for MR207827L4

Mdm2 (NM_010786) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Mdm2 (NM_010786) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Mdm2
Synonyms:	1700007J15Rik; AA415488; Mdm-2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR207827).
Restriction Sites:	Sgfl-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_010786
ORF Size:	1470 bp




[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_010786.2
RefSeq Size:	3124 bp
RefSeq ORF:	1470 bp
Locus ID:	17246
UniProt ID:	P23804
Cytogenetics:	10 66.32 cM
Gene Summary:	E3 ubiquitin-protein ligase that mediates ubiquitination of p53/TP53, leading to its degradation by the proteasome (PubMed:15195100, PubMed:21804542). Inhibits p53/TP53- and p73/TP73-mediated cell cycle arrest and apoptosis by binding its transcriptional activation domain (By similarity). Also acts as a ubiquitin ligase E3 toward itself, ARRB1 and ARRB2 (PubMed:11588219). Permits the nuclear export of p53/TP53 (By similarity). Promotes proteasome-dependent ubiquitin-independent degradation of retinoblastoma RB1 protein (By similarity). Inhibits DAXX-mediated apoptosis by inducing its ubiquitination and degradation (By similarity). Component of the TRIM28/KAP1-MDM2-p53/TP53 complex involved in stabilizing p53/TP53 (By similarity). Also component of the TRIM28/KAP1-ERBB4-MDM2 complex which links growth factor and DNA damage response pathways (By similarity). Mediates ubiquitination and subsequent proteasome degradation of DYRK2 in nucleus (By similarity). Ubiquitinates IGF1R and SNAI1 and promotes them to proteasomal degradation (By similarity). Ubiquitinates DCX, leading to DCX degradation and reduction of the dendritic spine density of olfactory bulb granule cells (PubMed:25088421). Ubiquitinates DLG4, leading to proteasomal degradation of DLG4 which is required for AMPA receptor endocytosis (PubMed:14642282).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR207827L4



Double digestion of MR207827L4 using SgfI and MluI