

Product datasheet for MR207859L3

Cyp2f2 (NM_007817) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Tag:	Myc-DDK
Symbol:	Cyp2f2
Synonyms:	Cyp2f
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR207859).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:




* The last codon before the Stop codon of the ORF.

ACCN:	NM_007817
ORF Size:	1473 bp



OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_007817.2 , NP_031843.2
RefSeq Size:	1842 bp
RefSeq ORF:	1476 bp
Locus ID:	13107
UniProt ID:	P33267
Cytogenetics:	7 A3
Gene Summary:	Involved in the regio- and stereoselective transformation of naphthalene to trans-1R-hydroxy-2R-glutathionyl-1,2-dihydronaphthalene in the presence of glutathione and glutathione S-transferases. It specifically catalyzes the production of a very reactive and potentially toxic intermediate, the 2R,2S arene oxide, that is associated with necrosis of the unciliated bronchiolar epithelial cells or Clara cells in lung.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR207859L3