

## Product datasheet for **MR207955L4V**

### Sp3 (BC027797) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Lentiviral Particles   |
| Product Name:             | Sp3 (BC027797) Mouse Tagged ORF Clone Lentiviral Particle  |
| Symbol:                   | Sp3  |
| Synonyms:                 | D130027J01Rik  |
| Mammalian Cell Selection: | Puromycin  |
| Vector:                   | pLenti-C-mGFP-P2A-Puro (PS100093)  |
| Tag:                      | mGFP   |
| ACCN:                     | BC027797   |
| ORF Size:                 | 1488 bp  |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(MR207955).   |
| OTI Disclaimer:           | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a> |
| OTI Annotation:           | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| RefSeq:                   | <a href="#">BC027797</a> , <a href="#">AAH27797</a>  |
| RefSeq Size:              | 3188 bp  |
| RefSeq ORF:               | 1490 bp  |
| Locus ID:                 | 20687  |
| Cytogenetics:             | 2 C3   |



[View online »](#)

**Gene Summary:**

This gene product belongs to a family of Sp1 related transcription factors, which regulate transcription by binding to consensus GC- and GT-box regulatory elements in target genes. This protein contains a zinc finger DNA-binding domain and several transactivation domains, and has been reported to function as a bifunctional transcription factor that either stimulates or represses transcription of numerous genes. Alternative splicing results in transcript variants encoding different isoforms, and one variant initiates translation from a non-AUG (AUA) codon. [provided by RefSeq, Jul 2008]