

Product datasheet for MR208041L3

Wasl (NM_028459) Mouse Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Taq: Myc-DDK

Symbol: Wasl

Synonyms: 2900021I12Rik; 3110031I02Rik; N-WASP

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)


E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide Sequence: The ORF insert of this clone is exactly the same as (MR208041).

Restriction Sites: Sgfl-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



The diagram illustrates the pET28b(+)-T7 promoter vector. It features a T7 promoter (T7P) with a transcription start site (ATG) indicated by a green arrow. The vector includes an *Eco*I site, a *Bam*H site, and an *RBS* (ribosome binding site) indicated by a blue arrow. A *Kozak Consensus* sequence is shown above the start site. A *Sgf*I site is located downstream. The vector ends with *Mlu*I, *Not*I, and *Xho*I sites, followed by a *Myc*.Tag sequence and a stop site (TAA).

T R T R P L E Q K L
DDK.Tag
T GAT ATC CTG GAT TAC AAG GAT GAC GAC GAT AAG GTT TGGGTAG

ACCN: NM 028459

ORE Size: 1503 bp




[View online »](#)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use.

©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_028459.2 , NP_082735.2
RefSeq Size:	4348 bp
RefSeq ORF:	1506 bp
Locus ID:	73178
UniProt ID:	Q9IYD9
Cytogenetics:	6 A3.1
Gene Summary:	Regulates actin polymerization by stimulating the actin-nucleating activity of the Arp2/3 complex. Involved in various processes, such as mitosis and cytokinesis, via its role in the regulation of actin polymerization. Together with CDC42, involved in the extension and maintenance of the formation of thin, actin-rich surface projections called filopodia. In addition to its role in the cytoplasm, also plays a role in the nucleus by regulating gene transcription, probably by promoting nuclear actin polymerization (By similarity). Binds to HSF1/HSTF1 and forms a complex on heat shock promoter elements (HSE) that negatively regulates HSP90 expression (PubMed:12871950). Plays a role in dendrite spine morphogenesis (PubMed:25851601). [UniProtKB/Swiss-Prot Function]

Product images:

Circular map for MR208041L3