

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for MR208216L4V

## Arcn1 (NM\_145985) Mouse Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Arcn1 (NM_145985) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Arcn1
Synonyms:	4632432M07Rik; nur17
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_145985
ORF Size:	1536 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR208216).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 145985.3, NP 666097.2</u>
RefSeq Size:	3470 bp
RefSeq ORF:	1536 bp
Locus ID:	213827
UniProt ID:	<u>Q5XJY5</u>
Cytogenetics:	9 A5.2



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:The coatomer is a cytosolic protein complex that binds to dilysine motifs and reversibly<br/>associates with Golgi non-clathrin-coated vesicles, which further mediate biosynthetic protein<br/>transport from the ER, via the Golgi up to the trans Golgi network. Coatomer complex is<br/>required for budding from Golgi membranes, and is essential for the retrograde Golgi-to-ER<br/>transport of dilysine-tagged proteins. In mammals, the coatomer can only be recruited by<br/>membranes associated to ADP-ribosylation factors (ARFs), which are small GTP-binding<br/>proteins; the complex also influences the Golgi structural integrity, as well as the processing,<br/>activity, and endocytic recycling of LDL receptors (By similarity).[UniProtKB/Swiss-Prot<br/>Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US