

Product datasheet for **MR208351L4V**

Snx1 (NM_019727) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Snx1 (NM_019727) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Snx1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_019727
ORF Size:	1566 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR208351).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_019727.2 , NP_062701.2
RefSeq Size:	2079 bp
RefSeq ORF:	1566 bp
Locus ID:	56440
UniProt ID:	Q9WV80
Cytogenetics:	9 C



[View online »](#)

Gene Summary:

Involved in several stages of intracellular trafficking. Interacts with membranes containing phosphatidylinositol 3-phosphate (PtdIns(3P)) or phosphatidylinositol 3,5-bisphosphate (PtdIns(3,5)P2). Acts in part as component of the retromer membrane-deforming SNX-BAR subcomplex. The SNX-BAR retromer mediates retrograde transport of cargo proteins from endosomes to the trans-Golgi network (TGN) and is involved in endosome-to-plasma membrane transport for cargo protein recycling. The SNX-BAR subcomplex functions to deform the donor membrane into a tubular profile called endosome-to-TGN transport carrier (ETC). Can sense membrane curvature and has in vitro vesicle-to-membrane remodeling activity. Involved in retrograde endosome-to-TGN transport of lysosomal enzyme receptors (IGF2R, M6PR and SORT1). Plays a role in targeting ligand-activated EGFR to the lysosomes for degradation after endocytosis from the cell surface and release from the Golgi. Involvement in retromer-independent endocytic trafficking of P2RY1 and lysosomal degradation of protease-activated receptor-1/F2R. Promotes KALRN- and RHO G-dependent but retromer-independent membrane remodeling such as lamellipodium formation; the function is dependent on GEF activity of KALRN. Required for endocytosis of DRD5 upon agonist stimulation but not for basal receptor trafficking (By similarity).[UniProtKB/Swiss-Prot Function]