

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR208599L4V

Kpna1 (NM_008465) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Kpna1 (NM_008465) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Kpna1
Synonyms:	AW494490; IPOA5; mSRP1; NPI1; Rch2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_008465
ORF Size:	1617 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR208599).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 008465.2</u>
RefSeq Size:	4022 bp
RefSeq ORF:	1617 bp
Locus ID:	16646
UniProt ID:	<u>Q60960</u>
Cytogenetics:	16 B3



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CRIGENEKpna1 (NM_008465) Mouse Tagged ORF Clone Lentiviral Particle - MR208599L4VGene Summary:Functions in nuclear protein import as an adapter protein for nuclear receptor KPNB1. Binds
specifically and directly to substrates containing either a simple or bipartite NLS motif.
Docking of the importin/substrate complex to the nuclear pore complex (NPC) is mediated by
KPNB1 through binding to nucleoporin FxFG repeats and the complex is subsequently
translocated through the pore by an energy requiring, Ran-dependent mechanism. At the
nucleoplasmic side of the NPC, Ran binds to importin-beta and the three components
separate and importin-alpha and -beta are re-exported from the nucleus to the cytoplasm
where GTP hydrolysis releases Ran from importin. The directionality of nuclear import is

Ran between the cytoplasm and nucleus.[UniProtKB/Swiss-Prot Function]

thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US