

## Product datasheet for **MR208633L4V**

### Grb10 (NM\_001177629) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Grb10 (NM_001177629) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Grb10
Synonyms:	5730571D09Rik; AI325020; Meg1; mKIAA0207
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001177629
ORF Size:	1626 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR208633).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_001177629.1</a> , <a href="#">NP_001171100.1</a>
RefSeq Size:	4764 bp
RefSeq ORF:	1626 bp
Locus ID:	14783
UniProt ID:	<a href="#">Q60760</a>
Cytogenetics:	11 7.15 cM



[View online »](#)

**Gene Summary:**

Adapter protein which modulates coupling of a number of cell surface receptor kinases with specific signaling pathways. Binds to, and suppress signals from, activated receptors tyrosine kinases, including the insulin (INSR) and insulin-like growth factor (IGF1R) receptors. The inhibitory effect can be achieved by 2 mechanisms: interference with the signaling pathway and increased receptor degradation. Delays and reduces AKT1 phosphorylation in response to insulin stimulation. Blocks association between INSR and IRS1 and IRS2 and prevents insulin-stimulated IRS1 and IRS2 tyrosine phosphorylation. Recruits NEDD4 to IGF1R, leading to IGF1R ubiquitination, increased internalization and degradation by both the proteasomal and lysosomal pathways. A similar role in the mediation of ubiquitination has also been suggested with INSR. Negatively regulates Wnt signaling by interacting with LRP6 intracellular portion and interfering with the binding of AXIN1 to LRP6. Positive regulator of the KDR/VEGFR-2 signaling pathway. May inhibit NEDD4-mediated degradation of KDR/VEGFR-2. [UniProtKB/Swiss-Prot Function]