

## Product datasheet for MR208654L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Papd7 (NM\_001169131) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Papd7 (NM\_001169131) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Papd7

Synonyms: LAK-1; POLK; Pols; TRF4; TRF4-1

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001169131

ORF Size: 1629 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR208654).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** NM 001169131.1, NP 001162602.1

 RefSeq Size:
 3994 bp

 RefSeq ORF:
 1626 bp

 Locus ID:
 210106

Cytogenetics: 13 35.55 cM







## **Gene Summary:**

Terminal nucleotidyltransferase that catalyzes preferentially the transfert of ATP and GTP on RNA 3' poly(A) tail creating a heterogeneous 3' poly(A) tail leading to mRNAs stabilization by protecting mRNAs from active deadenylation (By similarity). Also functions as a catalytic subunit of a TRAMP-like complex which has a poly(A) RNA polymerase activity and is involved in a post-transcriptional quality control mechanism. Polyadenylation with short oligo(A) tails is required for the degradative activity of the exosome on several of its nuclear RNA substrates. Has no terminal uridylyltransferase activity, and does not play a role in replication-dependent histone mRNA degradation via uridylation (By similarity).[UniProtKB/Swiss-Prot Function]