

Product datasheet for MR208663L3

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

Cyp1b1 (NM_009994) Mouse Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Cyp1b1 (NM_009994) Mouse Tagged Lenti ORF Clone

Tag: Myc-DDK
Symbol: Cyp1b1

Synonyms: CP1B; P4501b1

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(MR208663).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_009994

ORF Size: 1632 bp





Cyp1b1 (NM_009994) Mouse Tagged Lenti ORF Clone - MR208663L3

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 009994.1, NP 034124.1

RefSeq Size: 5128 bp RefSeq ORF: 1632 bp

Locus ID: 13078

UniProt ID: Q64429


Cytogenetics: 17 E3



Gene Summary:

A cytochrome P450 monooxygenase involved in the metabolism of various endogenous substrates, including fatty acids, steroid hormones and vitamins (By similarity). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (By similarity). Exhibits catalytic activity for the formation of hydroxyestrogens from 17beta-estradiol (E2), namely 2- and 4hydroxy E2 (PubMed:23821647). Metabolizes testosterone and progesterone to B or D ring hydroxylated metabolites (By similarity). May act as a major enzyme for all-trans retinoic acid biosynthesis in extrahepatic tissues. Catalyzes two successive oxidative transformation of alltrans retinol to all-trans retinal and then to the active form all-trans retinoic acid (PubMed:15258110). Catalyzes the epoxidation of double bonds of certain PUFA. Converts arachidonic acid toward epoxyeicosatrienoic acid (EpETrE) regioisomers, 8,9-, 11,12-, and 14,15- EpETrE, that function as lipid mediators in the vascular system (PubMed:15258110). Additionally, displays dehydratase activity toward oxygenated eicosanoids hydroperoxyeicosatetraenoates (HpETEs). This activity is independent of cytochrome P450 reductase, NADPH, and O2 (By similarity). Also involved in the oxidative metabolism of xenobiotics, particularly converting polycyclic aromatic hydrocarbons and heterocyclic aryl amines procarcinogens to DNA-damaging products (By similarity). Plays an important role in retinal vascular development. Under ambient/hyperoxic O2 conditions, promotes angiogenesis and capillary morphogenesis of retinal endothelial cells and pericytes, likely by metabolizing the oxygenated products symptomatic of oxidative stress (PubMed:19005183, PubMed:20032512, PubMed:23568032). Also, contributes to oxidative homeostasis and ultrastructural organization and function of trabecular meshwork tissue through modulation of POSTN expression (PubMed:23979599).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR208663L3