

## Product datasheet for **MR208673L3V**

### Cdyl (NM\_001123386) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Cdyl (NM_001123386) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Cdyl
Synonyms:	AI325931
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001123386
ORF Size:	1635 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR208673).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_001123386.1</a> , <a href="#">NP_001116858.1</a>
RefSeq Size:	3494 bp
RefSeq ORF:	1635 bp
Locus ID:	12593
UniProt ID:	<a href="#">Q9WTK2</a>
Cytogenetics:	13 14.39 cM



[View online »](#)

**Gene Summary:**

Isoform 2: Chromatin reader protein that recognizes and binds histone H3 trimethylated at 'Lys-9', dimethylated at 'Lys-27' and trimethylated at 'Lys-27' (H3K9me3, H3K27me2 and H3K27me3, respectively) (PubMed:12947414). Part of multimeric repressive chromatin complexes, where it is required for transmission and restoration of repressive histone marks, thereby preserving the epigenetic landscape (PubMed:12947414). Required for chromatin targeting and maximal enzymatic activity of Polycomb repressive complex 2 (PRC2); acts as a positive regulator of PRC2 activity by bridging the pre-existing histone H3K27me3 and newly recruited PRC2 on neighboring nucleosomes (By similarity). Acts as a corepressor for REST by facilitating histone-lysine N-methyltransferase EHMT2 recruitment and H3K9 dimethylation at REST target genes for repression (By similarity). Involved X chromosome inactivation in females: recruited to Xist RNA-coated X chromosome and facilitates propagation of H3K9me2 by anchoring EHMT2 (PubMed:24144980). Required for neuronal migration during brain development by repressing expression of RHOA (PubMed:28076783). In addition to act as a chromatin reader, acts as a hydro-lyase (By similarity). Shows crotonyl-coA hydratase activity by mediating the conversion of crotonyl-CoA ((2E)-butenoyl-CoA) to beta-hydroxybutyryl-CoA (3-hydroxybutanoyl-CoA), thereby acting as a negative regulator of histone crotonylation (By similarity). Histone crotonylation is required during spermatogenesis; down-regulation of histone crotonylation by CDYL regulates the reactivation of sex chromosome-linked genes in round spermatids and histone replacement in elongating spermatids (PubMed:28803779). May have histone acetyltransferase activity; such activity is however unsure in vivo (PubMed:12072557).[UniProtKB/Swiss-Prot Function]