

## Product datasheet for **MR208683L3V**

### Slc22a6 (NM\_008766) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Slc22a6 (NM_008766) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Slc22a6
Synonyms:	mOat1; NKT; Oat1; Orctl1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_008766
ORF Size:	1638 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR208683).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_008766.2</a>
RefSeq Size:	4026 bp
RefSeq ORF:	1638 bp
Locus ID:	18399
UniProt ID:	<a href="#">Q8VC69</a>
Cytogenetics:	19 5.44 cM



[View online »](#)

**Gene Summary:**

Involved in the renal elimination of endogenous and exogenous organic anions. Functions as organic anion exchanger when the uptake of one molecule of organic anion is coupled with an efflux of one molecule of endogenous dicarboxylic acid (glutarate, ketoglutarate, etc). Mediates the sodium-independent uptake of 2,3-dimercapto-1-propanesulfonic acid (DMPS), cidofovir, adefovir, 9-(2-phosphonylmethoxyethyl) guanine (PMEG), 9-(2-phosphonylmethoxyethyl) diaminopurine (PMEDAP), ochratoxin (OTA), acyclovir (ACV), 3'-azido-3'-deoxythymidine (AZT), cimetidine (CMD), 2,4-dichloro-phenoxyacetate (2,4-D), hippurate (HA), indoleacetate (IA), indoxyl sulfate (IS) and 3-carboxy-4-methyl-5-propyl-2-furanpropionate (CMPF) and edaravone sulfate (By similarity). Mediates the sodium-independent uptake of p-aminohippurate (PAH). PAH uptake is inhibited by benzothiazolylcysteine (BTC), S-chlorotrifluoroethylcysteine (CTFC), cysteine S-conjugates S-dichlorovinylcysteine (DCVC), furosemide, steviol, phorbol 12-myristate 13-acetate (PMA), calcium ionophore A23187, benzylpenicillin, bumetamide, losartan, probenecid, phenol red, urate, glutarate and alpha-ketoglutarate (By similarity). PAH uptake is inhibited by p-chloromercuribenzenesulphonate (PCMBS), diethyl pyrocarbonate (DEPC), indomethacin, sulindac, diclofenac, carprofen, okadaic acid and PKC activators.[UniProtKB/Swiss-Prot Function]