

Product datasheet for MR208703L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Trip10 (NM_134125) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Trip10 (NM 134125) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Trip10

Synonyms: Al646975; Cip4

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_134125 **ORF Size:** 1644 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR208703).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 134125.1</u>

 RefSeq Size:
 2290 bp

 RefSeq ORF:
 1644 bp

 Locus ID:
 106628

 UniProt ID:
 Q8C|53

Cytogenetics: 17 D







Gene Summary:

Required to coordinate membrane tubulation with reorganization of the actin cytoskeleton during endocytosis. Binds to lipids such as phosphatidylinositol 4,5-bisphosphate and phosphatidylserine and promotes membrane invagination and the formation of tubules. Also promotes CDC42-induced actin polymerization by recruiting WASL/N-WASP which in turn activates the Arp2/3 complex. Actin polymerization may promote the fission of membrane tubules to form endocytic vesicles. Required for the formation of podosomes, actin-rich adhesion structures specific to monocyte-derived cells. May be required for the lysosomal retention of FASLG/FASL (By similarity). Required for translocation of GLUT4 to the plasma membrane in response to insulin signaling.[UniProtKB/Swiss-Prot Function]