

Product datasheet for **MR208746L3V**

Cpsf6 (NM_001013391) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Cpsf6 (NM_001013391) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Cpsf6
Synonyms:	4733401N12Rik; AI256641; CFIM; CFIM68; HPBR11-4; HPBR11-7
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001013391
ORF Size:	1656 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR208746).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_001013391.1
RefSeq Size:	6534 bp
RefSeq ORF:	1656 bp
Locus ID:	432508
UniProt ID:	Q6NVF9
Cytogenetics:	10 D2


[View online »](#)

Gene Summary:

Component of the cleavage factor Im (CFIm) complex that functions as an activator of the pre-mRNA 3'-end cleavage and polyadenylation processing required for the maturation of pre-mRNA into functional mRNAs. CFIm contributes to the recruitment of multiprotein complexes on specific sequences on the pre-mRNA 3'-end, so called cleavage and polyadenylation signals (pA signals). Most pre-mRNAs contain multiple pA signals, resulting in alternative cleavage and polyadenylation (APA) producing mRNAs with variable 3'-end formation. The CFIm complex acts as a key regulator of cleavage and polyadenylation site choice during APA through its binding to 5'-UGUA-3' elements localized in the 3'-untranslated region (UTR) for a huge number of pre-mRNAs. CPSF6 enhances NUDT21/CPSF5 binding to 5'-UGUA-3' elements localized upstream of pA signals and promotes RNA looping, and hence activates directly the mRNA 3'-processing machinery. Plays a role in mRNA export. [UniProtKB/Swiss-Prot Function]