

# **Product datasheet for MR208772**

### Foxc1 (NM\_008592) Mouse Tagged ORF Clone

### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Foxc1 (NM\_008592) Mouse Tagged ORF Clone

Tag: Myc-DDK

Symbol: Foxc1

Synonyms: ch; fkh-1; Fkh1; FREAC3; frkhda; Mf1; Mf4

Mammalian Cell Neomycin

Selection:

**Vector:** pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





ORF Nucleotide Sequence:

>MR208772 representing NM\_008592
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ATGCAGGCGCGCTACTCGGTGTCCAGCCCCAACTCCCTGGGAGTGGTGCCCTACCTCGGCGGCGAGCAGA GCTACTATCGCGCTGCCGCGGCGGCCGGCCGGCGCCTACACCGCCATGCCGGCCCCTATGAGCGTGTA CTCGCACCCTGCTCACGCCGAGCAGTACCCGGGCAGCATGGCGCGCCCTACGGGCCTTATACGCCGCAG CCGCAGCCCAAGGACATGGTGAAGCCGCCCTACAGCTACATCGCTCTTATCACCATGGCCATCCAGAATG CCCCGGACAAGAAGATCACTCTGAATGGCATCTACCAGTTCATCATGGACCGCTTCCCCTTCTATCGGGA CAATAAGCAGGGCTGGCAGAACAGCATACGGCACAACCTCTCGCTCAACGAGTGCTTCGTCAAGGTGCCC CGCGACGACAAGAAGCCAGGCAAGGGCAGCTACTGGACGCTCGACCCGGACTCCTACAACATGTTCGAGA ACGGCAGCTTCCTGCGGCGGCGCGCGCTTCAAGAAGAAGGACGCAGTGAAGGACAAGGAGGAGAAGGA GCCGAGGGCTCCGCTCCCGGGCCACAGCCGCCGCCGTGCGCATCCAGGACATCAAGACGGAGAACGGTA CGTGTCCCTCGCCTCCCAGCCCCTGTCCCCGGCTGCCGCCCTAGGCAGCGGCAGCGCCACAGTGCC CAAAATCGAGAGCCCCGACAGCAGCAGCAGCTTGTCGAGCGGGAGCAGCCCCCGGGCAGCCTGCCG TCGGCGCGCCGCTCAGCCTGGACGCTGCAGAACCCGCGCCGCCGCCACAGCCTGCGCCGCCGCCGCATC ACAGCCAGGGCTTCAGCGTGGACAACATCATGACGTCGCTGCGGGGGTCGCCGCAGGGTTCGGCCGCTGA GCTCGGTTCCGGCCTCCTGGCCTCGGCGGCTGCGTCCTCGCGCGCGGGGCATCGCGCCCCCCGCTGGCGCTG GGTGCCTACTCTCCGGGCCAGAGCTCCCTCTACAGCTCCCCTGCAGCCAGAGCTCCAGTGCGGGCAGTT CGGGCGGCGGGGGTGGCGGCGGCGGAGGCGGCGGCAGCAGCAGCGCTGCGGGTACGGGGGGCGCCGC CCGGCGGGAGGCGCGGCGCGCGCGGCGGCGACCCCCCTGCCCGACTACTCGCTGCCTCCAGCGACCA GGGCCGACTCACCTCGTGGTACCTGAACCAGGCAGGTGGAGACCTGGGCCACTTGGCGAGCGCGGCGGCG GCTGCGGCGGCCGCAGGCTACCCTGGCCAGCAGCAGAACTTCCACTCGGTGCGGGAAATGTTCGAGTCTC AGCGGATCGGCTTGAACAACTCCCCGGTGAATGGGAATAGTAGCTGTCAGATGGCTTTCCCTGCCAGTCA GTCTCTGTACCGCACGTCGGGGGCTTTCGTCTATGACTGTAGCAAATTC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** 

>MR208772 representing NM\_008592 Red=Cloning site Green=Tags(s)

MQARYSVSSPNSLGVVPYLGGEQSYYRAAAAAAGGGYTAMPAPMSVYSHPAHAEQYPGSMARAYGPYTPQ PQPKDMVKPPYSYIALITMAIQNAPDKKITLNGIYQFIMDRFPFYRDNKQGWQNSIRHNLSLNECFVKVP RDDKKPGKGSYWTLDPDSYNMFENGSFLRRRRFKKKDAVKDKEEKGRLHLQEPPPPQAGRQPAPAPPEQ AEGSAPGPQPPPVRIQDIKTENGTCPSPPQPLSPAAALGSGSAATVPKIESPDSSSSLSSGSSPPGSLP SARPLSLDAAEPAPPPQPAPPPHHSQGFSVDNIMTSLRGSPQGSAAELGSGLLASAAASSRAGIAPPLAL GAYSPGQSSLYSSPCSQSSSAGSSGGGGGGGGGGGGSSSSAAGTGGAATYHCNLQAMSLYAAGERGGHLQG PAGGAGSAAVDDPLPDYSLPPATSSSSSSLSHGGGQEASHHPASHQGRLTSWYLNQAGGDLGHLASAAA AAAAAGYPGQQQNFHSVREMFESQRIGLNNSPVNGNSSCQMAFPASQSLYRTSGAFVYDCSKF


**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 


**Restriction Sites:** 

Sgfl-Mlul



**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_008592

ORF Size: 1659 bp

**OTI Disclaimer:** 

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customport@origene.com">customport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

**OTI Annotation:** 

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.



ORIGENE

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 008592.2</u>, <u>NP 032618.2</u>

 RefSeq Size:
 3990 bp

 RefSeq ORF:
 1662 bp

 Locus ID:
 17300

 UniProt ID:
 Q61572

**Cytogenetics:** 13 13.52 cM **MW:** 57.4 kDa

**Gene Summary:** DNA-binding transcriptional factor that plays a role in a broad range of cellular and

developmental processes such as eye, bones, cardiovascular, kidney and skin development

(PubMed:9635428, PubMed:9106663, PubMed:10479458, PubMed:10395790, PubMed:11562355, PubMed:18187037, PubMed:19668217, PubMed:22493429,

PubMed:24590069, PubMed:25808752, PubMed:28223138). Acts either as a transcriptional activator or repressor (PubMed:28223138). Binds to the consensus binding site 5'-[G/C] [A/T]AAA[T/C]AA[A/C]-3' in promoter of target genes (PubMed:25808752). Upon DNA-binding, promotes DNA bending. Acts as a transcriptional coactivator (PubMed:25808752). Stimulates

Indian hedgehog (Ihh)-induced target gene expression mediated by the transcription factor GLI2, and hence regulates endochondral ossification (PubMed:25808752). Acts also as a transcriptional coregulator by increasing DNA-binding capacity of GLI2 in breast cancer cells. Regulates FOXO1 through binding to a conserved element, 5'-GTAAACAAA-3' in its promoter region, implicating FOXC1 as an important regulator of cell viability and resistance to oxidative stress in the eye (By similarity). Cooperates with transcription factor FOXC2 in

regulating expression of genes that maintain podocyte integrity (PubMed:28223138). Promotes cell growth inhibition by stopping the cell cycle in the G1 phase through TGFB1-mediated signals. Involved in epithelial-mesenchymal transition (EMT) induction by increasing cell proliferation, migration and invasion (By similarity). Involved in chemokine CXCL12-

induced endothelial cell migration through the control of CXCR4 expression (PubMed:18187037). Plays a role in the gene regulatory network essential for epidermal keratinocyte terminal differentiation (By similarity). Essential developmental transcriptional factor required for mesoderm-derived tissues formation, such as the somites, skin, bone and cartilage (PubMed:9106663, PubMed:10479458, PubMed:10395790, PubMed:10704385, PubMed:11562355, PubMed:15196959). Positively regulates CXCL12 and stem cell factor expression in bone marrow mesenchymal progenitor cells, and hence plays a role in the


progenitor cells (HSPC) (PubMed:24590069). Plays a role in corneal transparency by preventing both blood vessel and lymphatic vessel growth during embryonic development in a VEGF-dependent manner (PubMed:22171010). May function as a tumor suppressor (By

development and maintenance of mesenchymal niches for haematopoietic stem and

similarity).[UniProtKB/Swiss-Prot Function]



## **Product images:**



Circular map for MR208772