

Product datasheet for **MR208794L3V**

Ubqln1 (NM_152234) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Ubqln1 (NM_152234) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Ubqln1
Synonyms:	1110046H03Rik; 1810030E05Rik; AU019746; C77538; D13ErtD372e; Da41; Dsk2; Plic-1; Plic1; Xdrp1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_152234
ORF Size:	1665 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR208794).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_152234.1 , NP_689420.1
RefSeq Size:	3609 bp
RefSeq ORF:	1665 bp
Locus ID:	56085
UniProt ID:	Q8R317
Cytogenetics:	13 30.95 cM



[View online »](#)

Gene Summary:

Plays an important role in the regulation of different protein degradation mechanisms and pathways including ubiquitin-proteasome system (UPS), autophagy and endoplasmic reticulum-associated protein degradation (ERAD) pathway. Mediates the proteasomal targeting of misfolded or accumulated proteins for degradation by binding (via UBA domain) to their polyubiquitin chains and by interacting (via ubiquitin-like domain) with the subunits of the proteasome. Plays a role in the ERAD pathway via its interaction with ER-localized proteins UBXLN4, VCP and HERPUD1 and may form a link between the polyubiquitinated ERAD substrates and the proteasome. Plays a role in unfolded protein response (UPR) by attenuating the induction of UPR-inducible genes, DDT13/CHOP, HSPA5 and PDIA2 during ER stress. Involved in the regulation of macroautophagy and autophagosome formation; required for maturation of autophagy-related protein LC3 from the cytosolic form LC3-I to the membrane-bound form LC3-II and may assist in the maturation of autophagosomes to autolysosomes by mediating autophagosome-lysosome fusion. Negatively regulates the TICAM1/TRIF-dependent toll-like receptor signaling pathway by decreasing the abundance of TICAM1 via the autophagic pathway. Plays a key role in the regulation of the levels of PSEN1 by targeting its accumulation to aggresomes which may then be removed from cells by autophagocytosis. Promotes the ubiquitination and lysosomal degradation of ORA11, consequently downregulating the ORA11-mediated Ca²⁺ mobilization. Suppresses the maturation and proteasomal degradation of amyloid beta A4 protein (A4) by stimulating the lysine 63 (K63)-linked polyubiquitination. Delays the maturation of A4 by sequestering it in the Golgi apparatus and preventing its transport to the cell surface for subsequent processing (By similarity). Links CD47 to the cytoskeleton (PubMed:10549293). [UniProtKB/Swiss-Prot Function]