

Product datasheet for **MR208916L3V**

Fbxw11 (NM_134015) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Symbol:	Fbxw11
Synonyms:	2310065A07Rik; AA536858; BTRC2; BTRCP2; Fbxw1b; HOS
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_134015
ORF Size:	1689 bp

ORF Nucleotide Sequence: The ORF insert of this clone is exactly the same as(MR208916).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

RefSeq:	NM_134015.2
RefSeq Size:	4339 bp
RefSeq ORF:	1692 bp
Locus ID:	103583
UniProt ID:	Q5SRY7
Cytogenetics:	11 A4



Gene Summary:

Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins. Probably recognizes and binds to phosphorylated target proteins. SCF(FBXW11) mediates the ubiquitination of phosphorylated CTNNB1 and participates in Wnt signaling. SCF(FBXW11) mediates the ubiquitination of phosphorylated NFKB1A, which degradation frees the associated NFKB1 to translocate into the nucleus and to activate transcription. SCF(FBXW11) mediates the ubiquitination of IFNAR1. SCF(FBXW11) mediates the ubiquitination of CEP68; this is required for centriole separation during mitosis (By similarity). Involved in the oxidative stress-induced a ubiquitin-mediated decrease in RCAN1. Mediates the degradation of CDC25A induced by ionizing radiation in cells progressing through S phase and thus may function in the intra-S-phase checkpoint. Has an essential role in the control of the clock-dependent transcription via degradation of phosphorylated PER1 and phosphorylated PER2. SCF(FBXW11) mediates the ubiquitination of CYTH1, and probably CYTH2 (By similarity).[UniProtKB/Swiss-Prot Function]