

Product datasheet for MR209113L3

UbqIn1 (NM_026842) Mouse Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids Product Name: Ubqln1 (NM_026842) Mouse Tagged Lenti ORF Clone Tag: Myc-DDK Symbol: Ubgln1 1110046H03Rik; 1810030E05Rik; AU019746; C77538; D13Ertd372e; Da41; Dsk2; Plic-1; Plic1; Synonyms: Xdrp1 Mammalian Cell Puromycin Selection: Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092) E. coli Selection: Chloramphenicol (34 ug/mL) **ORF** Nucleotide The ORF insert of this clone is exactly the same as(MR209113). Sequence: **Restriction Sites:** Sgfl-Mlul **Cloning Scheme:** Cloning sites used for ORF Shuttling: ORF Safl Mlu I

 $\begin{array}{c} \underbrace{Kozak}_{Consensus} \\ \underline{EcoR 1} \\ \underline{BamH1} \\ RBS \\ \underline{Sg1} \\ ORF \\ CTATAGGGCGGCCGGGAATTCGTCGACTGGATCGGTACCGAGGAGGATCTGCCGCGGATCGCC \\ ATG \\ \cdots \\ \cdots \\ nnn \\ \hline ACG \\ CGT \\ CTG \\ CGG \\ CGT \\ CTG \\ CGG \\ CGT \\ CTG \\ CGG \\$

--- GCG ATC GCC ATG --- // --- NNN ACG CGT ---

* The last codon before the Stop codon of the ORF.

ACCN: ORF Size: NM_026842 1746 bp OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


Service Content of the service of th	
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 026842.4</u>
RefSeq Size:	3693 bp
RefSeq ORF:	1749 bp
Locus ID:	56085
UniProt ID:	<u>Q8R317</u>
Cytogenetics:	13 30.95 cM

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE Ubqln1 (NM_026842) Mouse Tagged Lenti ORF Clone – MR209113L3

Plays an important role in the regulation of different protein degradation mechanisms and Gene Summary: pathways including ubiquitin-proteasome system (UPS), autophagy and endoplasmic reticulum-associated protein degradation (ERAD) pathway. Mediates the proteasomal targeting of misfolded or accumulated proteins for degradation by binding (via UBA domain) to their polyubiquitin chains and by interacting (via ubiquitin-like domain) with the subunits of the proteasome. Plays a role in the ERAD pathway via its interaction with ER-localized proteins UBXN4, VCP and HERPUD1 and may form a link between the polyubiguitinated ERAD substrates and the proteasome. Plays a role in unfolded protein response (UPR) by attenuating the induction of UPR-inducible genes, DDTI3/CHOP, HSPA5 and PDIA2 during ER stress. Involved in the regulation of macroautophagy and autophagosome formation; required for maturation of autophagy-related protein LC3 from the cytosolic form LC3-I to the membrane-bound form LC3-II and may assist in the maturation of autophagosomes to autolysosomes by mediating autophagosome-lysosome fusion. Negatively regulates the TICAM1/TRIF-dependent toll-like receptor signaling pathway by decreasing the abundance of TICAM1 via the autophagic pathway. Plays a key role in the regulation of the levels of PSEN1 by targeting its accumulation to aggresomes which may then be removed from cells by autophagocytosis. Promotes the ubiquitination and lysosomal degradation of ORAI1, consequently downregulating the ORAI1-mediated Ca2+ mobilization. Suppresses the maturation and proteasomal degradation of amyloid beta A4 protein (A4) by stimulating the lysine 63 (K63)-linked polyubiguitination. Delays the maturation of A4 by sequestering it in the Golgi apparatus and preventing its transport to the cell surface for subsequent processing (By similarity). Links CD47 to the cytoskeleton (PubMed:10549293).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR209113L3

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US