

# **Product datasheet for MR209117L3**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Ddi2 (BC031491) Mouse Tagged Lenti ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Ddi2 (BC031491) Mouse Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: Ddi2

**Synonyms:** 1110056G13Rik; 1700027M01Rik; 9130022E05Rik; Al604911; AU040698

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this

Sequence:

The ORF insert of this clone is exactly the same as(MR209117).

Restriction Sites: Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF.

**ACCN:** BC031491 **ORF Size:** 1746 bp



### Ddi2 (BC031491) Mouse Tagged Lenti ORF Clone - MR209117L3

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: BC031491

**RefSeq Size:** 2307 bp

RefSeq ORF: 1748 bp

**Locus ID:** 68817

Cytogenetics: 4 E1


Gene Summary: Aspartic protease that mediates the cleavage of NFE2L1/NRF1 at 'Leu-104', thereby promoting

release of NFE2L1/NRF1 from the endoplasmic reticulum membrane. Ubiquitination of NFE2L1/NRF1 is a prerequisite for cleavage, suggesting that DDI2 specifically recognizes and binds ubiquitinated NFE2L1/NRF1. Seems to act as a proteasomal shuttle which links the proteasome and replication fork proteins like RTF2. Required, with DDI1, for cellular survival following replication stress. Together or redudantly with DDI1, removes RTF2 from stalled forks to allow cell cycle progression after replication stress and maintains genome integrity.

[UniProtKB/Swiss-Prot Function]



# **Product images:**



Circular map for MR209117L3