

## Product datasheet for MR209117L3V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Ddi2 (BC031491) Mouse Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type: Lentiviral Particles

**Product Name:** Ddi2 (BC031491) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Ddi2

**Synonyms:** 1110056G13Rik; 1700027M01Rik; 9130022E05Rik; Al604911; AU040698

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: BC031491

ORF Size: 1746 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(MR209117).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

 RefSeq:
 BC031491

 RefSeq Size:
 2307 bp

 RefSeq ORF:
 1748 bp

 Locus ID:
 68817

 Cytogenetics:
 4 E1







#### **Gene Summary:**

Aspartic protease that mediates the cleavage of NFE2L1/NRF1 at 'Leu-104', thereby promoting release of NFE2L1/NRF1 from the endoplasmic reticulum membrane. Ubiquitination of NFE2L1/NRF1 is a prerequisite for cleavage, suggesting that DDI2 specifically recognizes and binds ubiquitinated NFE2L1/NRF1. Seems to act as a proteasomal shuttle which links the proteasome and replication fork proteins like RTF2. Required, with DDI1, for cellular survival following replication stress. Together or redudantly with DDI1, removes RTF2 from stalled forks to allow cell cycle progression after replication stress and maintains genome integrity. [UniProtKB/Swiss-Prot Function]