

Product datasheet for MR209132L4

Optn (NM_181848) Mouse Tagged Lenti ORF Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids						
Product Name:	Optn (NM_181848) Mouse Tagged Lenti ORF Clone						
Tag:	mGFP						
Symbol:	Optn						
Synonyms:	4930441O07Rik; FIP2; HYPL; NRP						
Mammalian Cell Selection:	Puromycin						
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)						
E. coli Selection:	Chloramphenicol (34 ug/mL)						
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR209132).						
Restriction Sites:	Sgfl-Mlul						
Cloning Scheme:							
	Cloning sites used for ORF Shuttling:						
	Sgf1 ORF Mlu I GCG ATC GC ATG// NNÑ ACG CGT						

<u>Ecor I</u> CTATAGGGCGGCCGGGAATTCGTC	 Bam GGA1		GTACC	RE	-	тста		 nsus gf I	- C C A	TG	 0	RF
NNŇ	 CGT R	ACG T		Vot I CCG P		GAG E	ATG M	GFP GGG G			 	- <u></u>
GGA CTC AGA GTT TGG GLRV * The last codon before the Stop c												

ACCN: ORF Size: NM_181848 1752 bp



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


	n (NM_181848) Mouse Tagged Lenti ORF Clone – MR209132L4
OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery.
	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Metho	 d: 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM 181848.4</u>
RefSeq Size:	2413 bp
RefSeq ORF:	1755 bp
Locus ID:	71648
UniProt ID:	<u>Q8K3K8</u>
Cytogenetics:	2 3.15 cM

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE Optn (NM_181848) Mouse Tagged Lenti ORF Clone – MR209132L4

Plays an important role in the maintenance of the Golgi complex, in membrane trafficking, in Gene Summary: exocytosis, through its interaction with myosin VI and Rab8. Links myosin VI to the Golgi complex and plays an important role in Golgi ribbon formation. Plays a role in the activation of innate immune response during viral infection. Mechanistically, recruits TBK1 at the Golgi apparatus, promoting its trans-phosphorylation after RLR or TLR3 stimulation. In turn, activated TBK1 phosphorylates its downstream partner IRF3 to produce IFN-beta. Plays a neuroprotective role in the eye and optic nerve. May act by regulating membrane trafficking and cellular morphogenesis via a complex that contains Rab8 and hungtingtin (HD). Mediates the interaction of Rab8 with the probable GTPase-activating protein TBC1D17 during Rab8mediated endocytic trafficking, such as of transferrin receptor (TFRC/TfR); regulates Rab8 recruitnment to tubules emanating from the endocytic recycling compartment. Autophagy receptor that interacts directly with both the cargo to become degraded and an autophagy modifier of the MAP1 LC3 family; targets ubiquitin-coated bacteria (xenophagy), such as cytoplasmic Salmonella enterica, and appears to function in the same pathway as SQSTM1 and CALCOCO2/NDP52. May constitute a cellular target for adenovirus E3 14.7, an inhibitor of TNF-alpha functions, thereby affecting cell death.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR209132L4

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US