

## Product datasheet for MR209176L4

### Hdac9 (NM\_024124) Mouse Tagged Lenti ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Hdac9 (NM_024124) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Hdac9
Synonyms:	AV022454; D030072B18Rik; HD7B; HD9; Hdac7b; HDRP; Mitr; mKIAA0744
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR209176).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


ACCN:	NM_024124
ORF Size:	1767 bp




[View online »](#)

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_024124.2</a> , <a href="#">NP_077038.2</a>
<b>RefSeq Size:</b>	4461 bp
<b>RefSeq ORF:</b>	1767 bp
<b>Locus ID:</b>	79221
<b>UniProt ID:</b>	<a href="#">Q99N13</a>
<b>Cytogenetics:</b>	12 A3
<b>Gene Summary:</b>	Devoided of intrinsic deacetylase activity, promotes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4) by recruiting HDAC1 and HDAC3. Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Represses MEF2-dependent transcription, inhibits skeletal myogenesis and may be involved in heart development. Protects neurons from apoptosis, both by inhibiting JUN phosphorylation by MAPK10 and by repressing JUN transcription via HDAC1 recruitment to JUN promoter. [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR209176L4



Double digestion of MR209176L4 using SgfI and MluI