

Product datasheet for MR209197L4

Trp73 (NM_001126330) Mouse Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids
Tag: mGFP
Symbol: Trp73
Synonyms: delta; deltaqNp73; p7; p73; TAp; TAp73; Tp73


Mammalian Cell: Puromycin
Selection:
Vector: pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide Sequence: The ORF insert of this clone is exactly the same as (MR209197).

Restriction Sites: Sgfl-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



The diagram shows the Kozak Consensus sequence: CTATAGGGCGGCCGGAAATTCTCGTCACTGGATCCGGTACCGAGGAGACTGCGCCGCGATCGC. Above the sequence, restriction sites are indicated: EcoRI, BamHI, RBS, SgfI, and MluI, NotI, XhoI, and mGFP Tag. The mGFP Tag is located at the 3' end. Below the sequence, the nucleotide sequence is shown: NNN (underlined), followed by ACG, CGT, ACG, CGG, CCG, CTC, GAG, ATG, AGC, GGG, GGC, and then a series of dashes. Below this, a second row shows the corresponding amino acids: G, L, R, V, P, L, E, M, S, G, G, and a series of dashes. The mGFP Tag is represented by a green box.

* The last codon before the Stop codon of the ORF.

ACCN: NM_001126330

ORF Size: 1770 bp



[View online »](#)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use.

©2025 OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, USA

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_001126330.1 , NP_001119802.1
RefSeq Size:	4756 bp
RefSeq ORF:	1773 bp
Locus ID:	22062
UniProt ID:	Q9JJP2
Cytogenetics:	4 83.79 cM

Gene Summary:

This gene encodes tumor protein p73, which is a member of the p53 family of transcription factors involved in cellular responses to stress and development. The family members include p53, p63, and p73 and have high sequence similarity to one another, which allows p63 and p73 to transactivate p53-responsive genes causing cell cycle arrest and apoptosis. The family members can interact with each other in many ways involving direct or indirect protein interactions, resulting in regulation of the same target gene promoters or regulation of each other's promoters. The p73 protein is expressed at very low levels in normal tissues and is differentially expressed in a number of tumors. The p73 gene expresses at least 35 mRNA variants due to the use of alternate promoters, alternate translation initiation sites, and multiple splice variations. Theoretically this can account for 29 different p73 isoforms; however, the biological validity and the full-length nature of most variants have not been determined. [provided by RefSeq, Jul 2008]

Product images: