

## Product datasheet for MR209236L3

### KIhI2 (NM\_178633) Mouse Tagged Lenti ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	KIhI2 (NM_178633) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	KIhI2
Synonyms:	6030411N21Rik; ABP-KELCH; AU020744; Mav
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR209236).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


ACCN:	NM_178633
ORF Size:	1782 bp



[View online >](#)

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_178633.2</a>
<b>RefSeq Size:</b>	3319 bp
<b>RefSeq ORF:</b>	1782 bp
<b>Locus ID:</b>	77113
<b>UniProt ID:</b>	<a href="#">Q8JZP3</a>
<b>Cytogenetics:</b>	8 B3.1
<b>Gene Summary:</b>	Promotes growth of cell projections in oligodendrocyte precursors. Responsible for degradative ubiquitination of the WNK kinases WNK1, WNK3 and WNK4 (By similarity). Component of a cullin-RING-based BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex that mediates the ubiquitination of target proteins, such as NPTXR, leading most often to their proteasomal degradation. Plays a role in the reorganization of the actin cytoskeleton. [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR209236L3