

## Product datasheet for MR209240L4V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Mtf2 (NM\_013827) Mouse Tagged ORF Clone Lentiviral Particle

### **Product data:**

Product Type: Lentiviral Particles

Product Name: Mtf2 (NM 013827) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Mtf2

**Synonyms:** 9230112N11Rik; AA537621; C76717; M96; Pcl2

**Mammalian Cell** 

- ·

Puromycin

Selection:

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_013827 **ORF Size:** 1779 bp

**ORF Nucleotide** 

- - -

Sequence:

The ORF insert of this clone is exactly the same as(MR209240).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 013827.3, NP 038855.2

RefSeq Size: 4343 bp
RefSeq ORF: 1782 bp
Locus ID: 17765
UniProt ID: Q02395

Cytogenetics: 5 F



#### **Gene Summary:**

Polycomb group (PcG) that specifically binds histone H3 trimethylated at 'Lys-36' (H3K36me3) and recruits the PRC2 complex. Acts by binding to H3K36me3, a mark for transcriptional activation, and recruiting the PRC2 complex, leading to enhance PRC2 H3K27me3 methylation activity. Regulates the transcriptional networks during embryonic stem cell self-renewal and differentiation. Promotes recruitment of the PRC2 complex to the inactive X chromosome in differentiating XX ES cells and PRC2 recruitment to target genes in undifferentiated ES cells. Required to repress Hox genes by enhancing H3K27me3 methylation of the PRC2 complex. In some conditions may act as an inhibitor of PRC2 activity: able to activate the CDKN2A gene and promote cellular senescence by suppressing the catalytic activity of the PRC2 complex locally. Binds to the metal-regulating-element (MRE) of MT1A gene promoter. [UniProtKB/Swiss-Prot Function]