

Product datasheet for MR209313L3

Arid3a (NM_007880) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Arid3a (NM_007880) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Arid3a
Synonyms:	Bright; Dri1; Drill
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR209313).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_007880
ORF Size:	1800 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_007880.2 , NP_031906.1
RefSeq Size:	5361 bp
RefSeq ORF:	1806 bp
Locus ID:	13496
UniProt ID:	Q62431
Cytogenetics:	10 C1
Gene Summary:	Transcription factor involved in B-cell differentiation. Binds a VH promoter proximal site necessary for induced mu-heavy-chain transcription. Binds the minor groove of a restricted ATC sequence that is sufficient for nuclear matrix association. This sequence motif is present in matrix-associating regions (MARS) proximal to the promoter and flanking E mu. Activates E mu-driven transcription by binding these sites. May be involved in the control of cell cycle progression by the RB1/E2F1 pathway.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR209313L3