

Product datasheet for MR209314L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cd96 (NM_032465) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Symbol: Cd96

Synonyms: 1700109112Rik; Tactile

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_032465

ORF Size: 1806 bp

ORF Nucleotide Sequence: The ORF insert of this clone is exactly the same as(MR209314).

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customercone care team at customercone contact our customer care team at customercone care team at <a href="mailto

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM_032465.1</u>, <u>NP_115854.1</u>





Cd96 (NM_032465) Mouse Tagged ORF Clone Lentiviral Particle | MR209314L3V

RefSeq Size: 2203 bp

RefSeq ORF: 1809 bp

Locus ID: 84544

UniProt ID: Q3U0X8

Cytogenetics: 16 B5

Gene Summary: May be involved in adhesive interactions of activated T and NK cells during the late phase of

the immune response. Promotes NK cell-target adhesion by interacting with PVR present on target cells. May function at a time after T and NK cells have penetrated the endothelium using integrins and selectins, when they are actively engaging diseased cells and moving

within areas of inflammation (By similarity).[UniProtKB/Swiss-Prot Function]