

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for MR209375L4V

## Galnt11 (NM\_144908) Mouse Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Galnt11 (NM_144908) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Galnt11
Synonyms:	A430075I06Rik; AI648252; E430002F06Rik
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_144908
ORF Size:	1827 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR209375).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 144908.3</u>
RefSeq Size:	2541 bp
RefSeq ORF:	1827 bp
Locus ID:	231050
UniProt ID:	<u>Q921L8</u>
Cytogenetics:	5 A3



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Polypeptide N-acetylgalactosaminyltransferase that catalyzes the initiation of protein O-<br/>linked glycosylation and is involved in left/right asymmetry by mediating O-glycosylation of<br/>NOTCH1. O-glycosylation of NOTCH1 promotes activation of NOTCH1, modulating the<br/>balance between motile and immotile (sensory) cilia at the left-right organiser (LRO).<br/>Polypeptide N-acetylgalactosaminyltransferases catalyze the transfer of an N-acetyl-D-<br/>galactosamine residue to a serine or threonine residue on the protein receptor. Displays the<br/>same enzyme activity toward MUC1, MUC4, and EA2 than GALNT1. Not involved in<br/>glycosylation of erythropoietin (EPO) (By similarity).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US