

## Product datasheet for **MR209644L4V**

### Cep290 (BC004690) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Cep290 (BC004690) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Cep290
Synonyms:	MGC7859
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	BC004690
ORF Size:	1907 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR209644).
OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:custsupport@origene.com">custsupport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a></p>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">BC004690</a> , <a href="#">AAH04690</a>
RefSeq Size:	2831 bp
RefSeq ORF:	1907 bp



[View online »](#)

Locus ID: 216274

Cytogenetics: 10 51.48 cM

**Gene Summary:** Involved in early and late steps in cilia formation (PubMed:21565611). Its association with CCP110 is required for inhibition of primary cilia formation by CCP110 (By similarity). May play a role in early ciliogenesis in the disappearance of centriolar satellites and in the transition of primary ciliar vesicles (PCVs) to capped ciliary vesicles (CCVs). Required for the centrosomal recruitment of RAB8A and for the targeting of centriole satellite proteins to centrosomes such as of PCM1 (By similarity). Required for the correct localization of ciliary and phototransduction proteins in retinal photoreceptor cells; may play a role in ciliary transport processes (PubMed:16632484). Required for efficient recruitment of RAB8A to primary cilium (By similarity). In the ciliary transition zone is part of the tectonic-like complex (also named B9 complex) which is required for tissue-specific ciliogenesis and may regulate ciliary membrane composition (PubMed:21725307). Involved in regulation of the BBSome complex integrity, specifically for presence of BBS2, BBS5 and BBS8/TTC8 in the complex, and in ciliary targeting of selected BBSome cargos. May play a role in controlling entry of the BBSome complex to cilia possibly implicating IQCB1/NPHP5 (By similarity). Activates ATF4-mediated transcription (By similarity).[UniProtKB/Swiss-Prot Function]