

### **Product datasheet for MR209692L3**

# Senp1 (NM\_144851) Mouse Tagged Lenti ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Senp1 (NM\_144851) Mouse Tagged Lenti ORF Clone

Tag: Myc-DDK
Symbol: Senp1

**Synonyms:** 2310046A20Rik; D15Ertd528e; E330036L07Rik; suPr-2

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(MR209692).

Sequence:

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF.

**ACCN:** NM\_144851

ORF Size: 1920 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

#### Senp1 (NM\_144851) Mouse Tagged Lenti ORF Clone - MR209692L3

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 144851.5</u>, <u>NP 659100.1</u>

 RefSeq Size:
 6415 bp

 RefSeq ORF:
 1923 bp

 Locus ID:
 223870

 UniProt ID:
 P59110

Cytogenetics: 15 54.04 cM

**Gene Summary:** Protease that catalyzes two essential functions in the SUMO pathway (PubMed:15923632,


PubMed:29499132). The first is the hydrolysis of an alpha-linked peptide bond at the C-terminal end of the small ubiquitin-like modifier (SUMO) propeptides, SUMO1, SUMO2 and SUMO3 leading to the mature form of the proteins. The second is the deconjugation of SUMO1, SUMO2 and SUMO3 from targeted proteins, by cleaving an epsilon-linked peptide bond between the C-terminal glycine of the mature SUMO and the lysine epsilon-amino group of the target protein. Deconjugates SUMO1 from HIPK2 (By similarity). Deconjugates SUMO1 from HDAC1 and BHLHE40/DEC1, which decreases its transcriptional repression activity (By similarity). Deconjugates SUMO1 from CLOCK, which decreases its transcriptional

activation activity (By similarity). Deconjugates SUMO2 from MTA1 (By similarity).

Deconjugates SUMO2 from MTA1 (By similarity). Deconjugates SUMO1 from METTL3 (By similarity). Desumoylates CCAR2 which decreases its interaction with SIRT1 (By similarity). Deconjugates SUMO1 from GPS2 (PubMed:29499132).[UniProtKB/Swiss-Prot Function]



## **Product images:**



Circular map for MR209692L3