

Product datasheet for MR209831L3

Wrnip1 (NM_030215) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Wrnip1 (NM_030215) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Wrnip1
Synonyms:	4833444L21Rik; WHIP; Wrnip
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR209831).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_030215
ORF Size:	1980 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_030215.3 , NP_084491.3
RefSeq Size:	2642 bp
RefSeq ORF:	1983 bp
Locus ID:	78903
UniProt ID:	Q91XU0
Cytogenetics:	13 A3.2
Gene Summary:	Functions as a modulator of initiation or reinitiation events during DNA polymerase delta-mediated DNA synthesis. In the presence of ATP, stimulation of DNA polymerase delta-mediated DNA synthesis is decreased. Plays also a role in the innate immune defense against viruses. Stabilizes the RIG-I/DDX58 dsRNA interaction and promotes RIG-I/DDX58 'Lys-63'-linked polyubiquitination. In turn, RIG-I/DDX58 transmits the signal through mitochondrial MAVS.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR209831L3