

Product datasheet for **MR209855L4V**

Appl2 (NM_145220) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Appl2 (NM_145220) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Appl2
Synonyms:	Dip3b
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_145220
ORF Size:	1989 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR209855).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_145220.1
RefSeq Size:	2958 bp
RefSeq ORF:	1989 bp
Locus ID:	216190
UniProt ID:	Q8K3G9
Cytogenetics:	10 C1



[View online »](#)

Gene Summary:

Multifunctional adapter protein that binds to various membrane receptors, nuclear factors and signaling proteins to regulate many processes, such as cell proliferation, immune response, endosomal trafficking and cell metabolism (PubMed:25568335, PubMed:27219021, PubMed:25328665, PubMed:19661063, PubMed:29467283). Regulates signaling pathway leading to cell proliferation through interaction with RAB5A and subunits of the NuRD/MeCP1 complex (By similarity). Plays a role in immune response by modulating phagocytosis, inflammatory and innate immune responses (PubMed:25568335, PubMed:27219021, PubMed:25328665). In macrophages, enhances Fc-gamma receptor-mediated phagocytosis through interaction with RAB31 leading to activation of PI3K/Akt signaling (PubMed:25568335). In response to LPS, modulates inflammatory responses by playing a key role on the regulation of TLR4 signaling and in the nuclear translocation of RELA/NF-kappa-B p65 and the secretion of pro- and anti-inflammatory cytokines (PubMed:27219021). Also functions as a negative regulator of innate immune response via inhibition of AKT1 signaling pathway by forming a complex with APPL1 and PIK3R1 (PubMed:25328665). Plays a role in endosomal trafficking of TGFBR1 from the endosomes to the nucleus (By similarity). plays a role in cell metabolism by regulating adiponectin and insulin signaling pathways and adaptive thermogenesis (PubMed:19661063, PubMed:29467283) (By similarity). In muscle, negatively regulates adiponectin-stimulated glucose uptake and fatty acid oxidation by inhibiting adiponectin signaling pathway through APPL1 sequestration thereby antagonizing APPL1 action (PubMed:19661063). In muscles, negatively regulates insulin-induced plasma membrane recruitment of GLUT4 and glucose uptake through interaction with TBC1D1 (By similarity). Plays a role in cold and diet-induced adaptive thermogenesis by activating ventromedial hypothalamus (VMH) neurons through AMPK inhibition which enhances sympathetic outflow to subcutaneous white adipose tissue (sWAT), sWAT being and cold tolerance (PubMed:29467283). Also plays a role in other signaling pathways namely Wnt/beta-catenin, HGF and glucocorticoid receptor signaling (PubMed:28965332, PubMed:29675572, PubMed:26445298). Positive regulator of beta-catenin/TCF-dependent transcription through direct interaction with RUVBL2/reptin resulting in the relief of RUVBL2-mediated repression of beta-catenin/TCF target genes by modulating the interactions within the beta-catenin-reptin-HDAC complex (By similarity). May affect adult neurogenesis in hippocampus and olfactory system via regulating the sensitivity of glucocorticoid receptor (PubMed:28965332, PubMed:29675572). Required for fibroblast migration through HGF cell signaling (PubMed:26445298).[UniProtKB/Swiss-Prot Function]