

Product datasheet for **MR209866L3V**

Chfr (NM_172717) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Chfr (NM_172717) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Chfr
Synonyms:	5730484M20Rik; C230082M18; RNF116
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_172717
ORF Size:	1989 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR209866).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_172717.3
RefSeq Size:	3166 bp
RefSeq ORF:	1992 bp
Locus ID:	231600
UniProt ID:	Q810L3
Cytogenetics:	5 F



[View online »](#)

Gene Summary:

E3 ubiquitin-protein ligase that functions in the antephasis checkpoint by actively delaying passage into mitosis in response to microtubule poisons. Acts in early prophase before chromosome condensation, when the centrosomes move apart from each other along the periphery of the nucleus. Probably involved in signaling the presence of mitotic stress caused by microtubule poisons by mediating the 'Lys-48'-linked ubiquitination of target proteins, leading to their degradation by the proteasome. Promotes the ubiquitination and subsequent degradation of AURKA and PLK1. Probably acts as a tumor suppressor, possibly by mediating the polyubiquitination of HDAC1, leading to its degradation. May also promote the formation of 'Lys-63'-linked polyubiquitin chains and functions with the specific ubiquitin-conjugating UBC13-MMS2 (UBE2N-UBE2V2) heterodimer. Substrates that are polyubiquitinated at 'Lys-63' are usually not targeted for degradation, but are rather involved in signaling cellular stress (By similarity).[UniProtKB/Swiss-Prot Function]