

Product datasheet for **MR209896**

Leo1 (NM_001039522) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Leo1 (NM_001039522) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Leo1
Synonyms:	Gm185
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

**ORF Nucleotide
Sequence:**

>MR209896 ORF sequence
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGGCGGACATGGAGACCTCTTCGGGAGCGAAGCTGAGAGCGAGGCCGAGCGCAAAGATTCCGAGTCTG
 AGTCCGACTCGGACTCTGACCAGGACAACGGCGCTTCTGGCAGCAACGCCTCTGGCAGCGAAAGTGATCA
 GGATGACAGAGGTGACTCCGGACAGCCAGTAACAAGGAAGTCTCGGGACGACAGTGAGGAGGAGGGG
 GCCTCTCATCACAGTGGCAGTGACAATCACTCTGAACGGTCAGACAATAGGTCGGAGGCTTCCGAGCGCT
 CTGACCACGAGGATAATGAGCCCTCTGACGAGGACCAGCACAGTGGCTCTGAAGCCACAACGATGACGA
 CGATGAGGGCCACAGGTCGGACGAAGGGAGCCGTCCTCAGAGGCAGAAGGCTCTGAAAAAGCCAGTCA
 GATGATGAGAAAATGGGATGGAGAGGATAAAAGTGACCAGTCAGATGACGAGAAGTGCAGAACTCTGACG
 ACGAGGATCGGGAGCAAGGTCGGATGAGGATAAGCTGCAGAATCTGACGACGACGAGGAGAAAATGCA
 GAACACAGATGACGAGGACCGGGCCAGATTTCTGATGATGACAGACAGCAGCTGTCTGAGGAGGAGAAG
 GGGAAATCTGACGATGAGCATCCAGTAGCTTCTGATAATGACGAGGAGAAGCAGAATTCGGACGATGAGG
 ACCAGCCCGAGGTGTCCGATGAGGAGAAAATGCAGAACTCAGACGATGAAAGGCCACAAGTCTCAGATGA
 AGATGGAAGGCGTTCAGACGGTGGAGGAGCAAGACCAGAAGTCAGAATCTGCAAGAGGCGAGTGACAGT
 GAAGATGAAGTTTTACGACTGAAGCGCAAGAACCGGATTCATCCGATTCCGAAGCAGACAGCGACACCG
 AGGTGCCCAAAGATAATAATGGAACCATGGATCTGTTCCGAGGTGCAGATGACATATCTTCAGGGAGTGA
 TGGAGAAGATAAGCCCCAACTCCAGGACAGCCTGTGGATGAAAATGGCTTGCCCCAGGATCAGCAGGAG
 GAAGAGCCCATACCTGAGACCAGAATAGAAGTGGAGATCCCCAAAGTAAACACGGACTTGGGCAATGACT
 TATATTTTGTAAACTGCCCAACTTCTCAGTGTAGAGCCAGGCCTTTTGATCCTCAGTATTATGAAGA
 TGAATTTGAAGATGAGGAGATGCTGGATGAAGAGGGGAGAACCAGGTTAAAATTGAAGGTAGAAAACT
 ATAAGATGGAGGATACGCAGAGATGAAGAAGGAAATGAAATTAAGAAAAGCAATGCGCGGATCGTCAAGT
 GGTGAGATGGAAGCATGTCCCTGCATTTAGGGAATGAAGTGTTTGATGTGTACAAAGCCCCACTGCAGGG
 CGACCACAACCATCTTTTCATAAGACAAGTACTGGGCTGCAGGGGCAGGCCGCTTTTAAACGAAGCTC
 ACCTTCAGGCCTCACTCTACAGACAGTGACACACAGGAAGATGACCCTGTCACTTGCAGATAGATGCT
 CAAAGACACAGAAGATTCGAATCTTACCAATGGCTGGCCGTGACCCTGAGTGCCAGCGCACAGAGATGAT
 TAAGAAAAGAAGAAGAACGCTGAGGGCCTCCATCAGGAGGGAGTCTCAGCAGCGCCGGATGCGAGAGAAA
 CAGCACCAGCGGGGGCTGAGCGCCAGCTACCTGGAGCCTGATCGTACGACGAGGAGGAGGAAGGCGAGG
 AGTCCGTCAGCCTGGCTGCCATTA AAAACCGCTACAAAGGGGGCATCCGCGAGGAACGAGCCAGAATCTA
 CTCTCGGACAGTGATGAGGGCTCCGAAGAAGCAAAAGCTCAAAGGTTACTCAAAGCAAAGAAGCTCAAC
 AGTGATGAGGAAGGTGAATCTTCTGAAAAGAGGAAAGCGGAAGACGATGACAAAGCGAATAAAAAGCATA
 AGAAGTATGTGATCAGTATGAAGAAGAAGAAGATGAC

ACGCGTACGCGGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR209896 protein sequence
Red=Cloning site Green=Tags(s)

MADMEDLFGSEAESEAERK DSESESDSDSDQDNGASGSNASGSESDQDDRGDSGQPSNKELFGDDSEEEG
 ASHHSGSDNHSESRSDNRSEASERSDHEDNEP SDEDQHSGSEAHNDDDDGHRSDGSRHSEAEGSEKAQS
 DDEKWDGEDKSDQSDDEKLQNSDDEDREQGSDEDKLQNSDDDEEKMQNTDDEDRAQISDDDRQQLSEEEK
 GNSDDEHPVASDNDEEKQNSDDEDQPQVSDDEEKMQNSDDERPQVSDDEDGRRSDGEEEQDQKSEARGSDS
 EDEVLR LKRKNAIPSDSEADSDTEVPKDNNGTMDLFGGADDISSGSDGEDKPPTPGQPVDENGLPQDQQE
 EEP I PETRIEVEIPKVNTDLGNDLYFVKLPNFLSVEPRPFDPPYYEDEFDEEMLDEEGRTRLKLVKVENT
 IRWRIRRDEEGNEIKESNARIVKWSGMSLHLGNEVFDVYKAPLQGDHNLH FIRQGTGLQGQAVFKTKL
 TFRPHSTSDTHRKM TSLADRC SKTQKIRILPMAGRDP ECQRTEMIKKEEERLRASIRRESQQRMRREK
 QHQ RGLSASYLEPDRYDEEEEGEESVSLAAIKNRYKGGIREERARIYSSDSDEGSEEDKAQRLLKAKKLN
 SDEEGESSGKRKAEDDDKANKKHKYV I SDEEEEDD

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:



ACCN: NM_001039522

ORF Size: 2004 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001039522.1](#), [NP_001034611.1](#)

RefSeq Size: 2192 bp

RefSeq ORF: 2004 bp

Locus ID: 235497


UniProt ID: [Q5XJE5](#)

Cytogenetics: 9 D

MW: 75.6 kDa

Gene Summary: Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser-5'-phosphorylated forms and is involved in transcriptional elongation, acting both independently and synergistically with TCEA1 and in cooperation with the DSIF complex and HTATSF1. PAF1C is required for transcription of Hox and Wnt target genes. PAF1C is involved in hematopoiesis and stimulates transcriptional activity of KMT2A/MLL1. PAF1C is involved in histone modifications such as ubiquitination of histone H2B and methylation on histone H3 'Lys-4' (H3K4me3). PAF1C recruits the RNF20/40 E3 ubiquitin-protein ligase complex and the E2 enzyme UBE2A or UBE2B to chromatin which mediate monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1); UB2A/B-mediated H2B ubiquitination is proposed to be coupled to transcription. PAF1C is involved in mRNA 3' end formation probably through association with cleavage and poly(A) factors. Involved in polyadenylation of mRNA precursors. Connects PAF1C to Wnt signaling (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR209896