

Product datasheet for **MR209902L3V**

Kcnq1 (NM_008434) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Kcnq1 (NM_008434) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Kcnq1
Synonyms:	AW559127; Kcna9; KVLQT1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_008434
ORF Size:	2007 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR209902).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_008434.2 , NP_032460.2
RefSeq Size:	3052 bp
RefSeq ORF:	2007 bp
Locus ID:	16535
UniProt ID:	P97414
Cytogenetics:	7 88.12 cM



[View online »](#)

Gene Summary:

Potassium channel that plays an important role in a number of tissues, including heart, inner ear, stomach and colon (By similarity) (PubMed:16314573, PubMed:11120752, PubMed:15004216). Associates with KCNE beta subunits that modulates current kinetics (By similarity) (PubMed:17597584, PubMed:15004216). Induces a voltage-dependent by rapidly activating and slowly deactivating potassium-selective outward current (By similarity) (PubMed:8900282). Promotes also a delayed voltage activated potassium current showing outward rectification characteristic (By similarity). During beta-adrenergic receptor stimulation participates in cardiac repolarization by associating with KCNE1 to form the I(Ks) cardiac potassium current that increases the amplitude and slows down the activation kinetics of outward potassium current I(Ks) (By similarity) (PubMed:15004216, PubMed:17597584). Muscarinic agonist oxotremorine-M strongly suppresses KCNQ1/KCNE1 current (By similarity). When associated with KCNE3, forms the potassium channel that is important for cyclic AMP-stimulated intestinal secretion of chloride ions (By similarity). This interaction with KCNE3 is reduced by 17beta-estradiol, resulting in the reduction of currents (By similarity). During conditions of increased substrate load, maintains the driving force for proximal tubular and intestinal sodium ions absorption, gastric acid secretion, and cAMP-induced jejunal chloride ions secretion (PubMed:16314573). Allows the provision of potassium ions to the luminal membrane of the secretory canaliculus in the resting state as well as during stimulated acid secretion (PubMed:19491250). When associated with KCNE2, forms an heterooligomer complex leading to currents with an apparently instantaneous activation, a rapid deactivation process and a linear current-voltage relationship and decreases the amplitude of the outward current (By similarity). When associated with KCNE4, inhibits voltage-gated potassium channel activity (By similarity). When associated with KCNE5, this complex only conducts current upon strong and continued depolarization (By similarity). Also forms a heterotetramer with KCNQ5; has a voltage-gated potassium channel activity (By similarity). Binds with phosphatidylinositol 4,5-bisphosphate (By similarity).[UniProtKB/Swiss-Prot Function]