

Product datasheet for **MR210449**

Pla2g4a (NM_008869) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Pla2g4a (NM_008869) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Pla2g4a
Synonyms:	cP; cPL; cPLA2; cPLA2-alpha; cPLA2alpha; Pla; Pla2g4
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide
Sequence:

>MR210449 representing NM_008869
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGCATCGCC**

ATGCTTTTCATAGATCCTTATCAGCACATTATAGTGAACACCAGTACTCCATAAGTTTACTGTTGTGG
TTCTACGTGCCACCAAAGTAACCAAGGGGACCTTTGGCGATATGCTGGACACTCCAGATCCTTATGTGGA
ACTTTTCATCTCTACAACCCCTGACAGCAGGAAGCGAACGAGACACTTCAATAATGATATAAACCCCGTG
TGGAAATGAGACCTTTGAGTTCATTTTGGATCCTAATCAGGAAAATGTTTTGGAGATCACACTGATGGATG
CCAACACTACGTATGGATGAAACCCTAGGCACAGCTACATTCCTGTATCTTCAATGAAAGTGGGAGAGAA
GAAAGAAGTCCTTTTATTTCAACCAAGTCACTGAAATGATTCTGAAAATGTCTCTGGAAGTTTGTTC
TGCCAGACCTACGGTTCAGCATGGCACTGTGTGATCAGGAGAAAACCTTCAGACAGCAGAGGAAAGAGA
ACATAAAAGAGAACATGAAGAACTTTTGGTCCAAAAAAGAGTGAGGGGCTTTATCCACACGTGATGT
GCCGGTGGTGGCCATTTTGGGTTCAAGTGGGGTTTCCGGGCCATGGTGGGATTCTCTGGTGTGATGAAG
GCACTGTATGAGTCGGGGATTTTGGACTGTGCTACATACATTGCTGGTCTTTCTGGATCCACATGGTACA
TGCAACCTTGTACTCTACCCCGATTTTCCAGAGAAAAGTCCCAGGAGATTAATGAAGAGCTAATGAA
AAATGTGACGCCACAACCTCTCTTACTTCTTACACCACAGAAAAGTAAAAGATACGTTGAGTCTTTATGG
AAGAAGAAAAGTTCTGGCCAGCCTGTCACCTTTACTGACATCTTTGGGATGCTAATAGGAGAAAACCTAA
TTCAAAATAGGATGAGCATGACCCTGAGTAGTTGAAGGAAAAGGTCAATGCCGCCCGGTGTCCTTTGCC
TCTCTTACGTTCTCCACGTCAAACCTGATGTGTGAGAGCTGATGTTTGGCGATTGGGTGGAATTTAGT
CCATATGAGATTGGCATGGCAAAATATGGTACCTTTATGGCTCCTGACCTATTTGGAAGCAAGTTTTTTA
TGGGAACAGTTGTAATAAATATGAAGAAAACCCCTTGCATTTCTTGATGGGTGTCTGGGCAGTGCCTT
TTCTATACTGTTCAACAGAGTTTTGGGAGTTTCTGGCTCACAGAATAAAGGCTCTACAATGGAAGAGGAA
TTAGAAAATATTACAGCAAAGCACATCGTGAGTAATGACAGCTCCGACAGTGATGATGAGGCTCAAGGAC
CCAAAGGCACCGAGAATGAAGAAGCTGAAAAAGAGTACCAAAGCGACAACCAAGCAAGTTGGGTCCATCG
GATGCTAATGGCCTTGGTGGGACTCGGCTTTATTCAATACCCGAGAAGGACGTGCCGAAAGGTGCAT
AACTTCATGCTGGGCTTGAATCTCAACACATCATATCCACTGTCTCCCCTGAGAGACTTCAGCTCTCAGG
ATTCTTCGATGACGAGCTCGACGACGCGGTAGCAGATCCAGATGAATTTGAACGAATATATGAACCACT
GGATGTCAAAAGTAAGAAGATTCATGTGGTAGATAGTGGGCTCACATTTAACCTGCCATATCCCTTGATT
CTTCGACCTCAGAGAGGTGTGGATCTTATCATCTCCTTTGACTTTCTGCAAGGCCGAGTGACACCAGTC
CCCTTTCAAGGAACTTCTGCTTGCAGAGAAGTGGCGAAAATGAACAAGCTTCCCTTTCCAAAGATCGA
TCCTTATGTGTTTATGATCGGGAAGGATTAAGGAATGCTATGTTTTAAACCTAAGAACTCCTGATGTGGAG
AAGGATTGCCCAACCATATCCACTTTGTTCTGGCCAACATCAACTTCAGAAAAGTACAAGGCCCCAGGTG
TTCTAAGGAAAACCAAAGAAGAGAAAGAAATGCTGACTTTGACATTTTGGATGACCCCGAATCGCCATT
TTCAACCTTCAACTTTTCAATATCCCAATCAAGCATTCAAAGGCTTCACGATTTGATGACTTCAACACA
CTGAACAACATTGATGTGATAAAGGATGCCATTTGTTGAGAGCATTGAATACAGAAGACAGAACCCATCTC
GTTGCTCTGTTTCCCTCAGTAATGTTGAAGCAAGAAAATCTTCAATAAGGAGTTTCAAGTAAACCCAC
TGTG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR210449 representing NM_008869
 Red=Cloning site Green=Tags(s)

```
MSFIDPYQHIIVEHQYSHKFTVVVLRATKVTKGTFGDMLDTPDPYVELFISTTPDSRKRTRHFNNDINPV
WNETFEFILDPNQENVLEITLMDANYVMDELTGTATFPVSSMKVGEKKEVPVIFNQVTEMIEMSLEVCS
CPDLRFSMALCDQEKTFRQQRKENIKENMKLLGPKKSEGLYSTRDVPVVAIILGSGGGFRAMVGFSGVMK
ALYESGILDCAIYIAGLSGSTWYMSTLYSHPDFPEKGPEEINEELMKNVSHNPLLLLTPQKVKRYVESLW
KKKSSGQPVTFTDIFGMLIGETLIQNRMSMTLSSLKEKVNAARCPLPLFTCLHVKPDVSELMFADWVEFS
PYEIGMAKYGTFFMAPDLFGSKFFMGTVVKKYEENPLHFLMGVWGSFAFSLFNRVLGVSGSQKNGSTMEEE
LENITAKHIVSNDSSDDEAQGPKGTENEEAEKEYQSDNQASWVHRMLMALVSDSALFNTREGRAGKVH
NFMLGLNLNTSYPLSPLRDFSSQDSFDELDAVADPDEFERIEPLDVKSKIHVVDSGLTFNLPYPLI
LRPQRGVDLIISFDFSARPSDTSPPFKELLLAEKWAKMNLPPFKIDPVYFDREGLKECYVFKPNPDVE
KDCPTIIHFVLANINFRKYKAPGVLRETKEEKEIADFDIFDDPESPFSTFNFQYPNQAFKRLHDLMYFNT
LNNIDVIKDAIVESIEYRRQNPSCRCSVLSNVEARKFFNKEFLSKPTV
```


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mm9034_h06.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_008869


ORF Size: 2244 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_008869.4
RefSeq Size:	2846 bp
RefSeq ORF:	2247 bp
Locus ID:	18783
UniProt ID:	P47713
Cytogenetics:	1 63.51 cM
MW:	85.7 kDa
Gene Summary:	<p>The protein encoded by this gene is a member of the phospholipase A2 group IV family. This enzyme hydrolyzes membrane phospholipids, thereby releasing the polyunsaturated fatty acid, arachidonic acid. Arachidonic acid is further metabolized into eicosanoids such as leukotrienes, thromboxanes and prostaglandins, that play important roles in regulating diverse biological processes such as inflammatory responses, membrane and actin dynamics, and tumorigenesis. A rise in intracellular calcium levels results in binding of calcium to the C2 domain of this protein, and triggers the translocation from the cytosol to intracellular membranes, including the Golgi apparatus. Disruption of this gene in mice led to decreased levels of eicosanoids and platelet-activating factor, decreased allergic symptoms, and impaired reproductive ability in females. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Mar 2015]</p>

Product images:



Circular map for MR210449