


Plasmid Map:


ACCN: NM_007970

ORF Size: 2253 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_007970.2](#), [NP_031996.1](#)

RefSeq Size: 4224 bp

RefSeq ORF: 2253 bp

Locus ID: 14055

UniProt ID: [P70351](#)

Cytogenetics: 11 64.37 cM

Gene Summary: This gene encodes a member of the Polycomb-group (PCG) family. The encoded protein is interchangeable with the related Enhancer of zeste 2 (Ezh2) protein as a core component of the polycomb repressive complex 2 (PRC2), which methylates histone H3 at lysine 27 and results in the transcriptional repression of affected target genes. This complex is involved in carrying out cell-fate decisions during embryonic stem cell differentiation. [provided by RefSeq, Sep 2014]