

## Product datasheet for **MR210487L3V**

### Large1 (NM\_010687) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Large1 (NM_010687) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Large1
Synonyms:	BPFD#36; enr; fg; froggy; Gyltl1a; Large; Mbp-1; Mbp1; myd
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_010687
ORF Size:	2268 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR210487).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_010687.1</a> , <a href="#">NP_034817.1</a>
RefSeq Size:	3669 bp
RefSeq ORF:	2271 bp
Locus ID:	16795
UniProt ID:	<a href="#">Q9Z1M7</a>
Cytogenetics:	8 35.08 cM



[View online »](#)

**Gene Summary:**

Bifunctional glycosyltransferase with both xylosyltransferase and beta-1,3-glucuronyltransferase activities involved in the biosynthesis of the phosphorylated O-mannosyl trisaccharide (N-acetylgalactosamine-beta-3-N-acetylglucosamine-beta-4-(phosphate-6-)mannose), a carbohydrate structure present in alpha-dystroglycan (DAG1) (PubMed:23125099, PubMed:23135544). Phosphorylated O-mannosyl trisaccharid is required for binding laminin G-like domain-containing extracellular proteins with high affinity and plays a key role in skeletal muscle function and regeneration (PubMed:15184894, PubMed:24132234). LARGE elongates the glucuronyl-beta-1,4-xylose-beta disaccharide primer structure initiated by B3GNT1/B4GAT1 by adding repeating units [-3-Xylose-alpha-1,3-GlcA-beta-1-] to produce a heteropolysaccharide (By similarity).[UniProtKB/Swiss-Prot Function]