

Product datasheet for **MR210500**

Ikbbk (NM_001159774) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Ikbbk (NM_001159774) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Ikbbk
Synonyms:	A1132552; IKK-2; IKK-beta; IKK2; IKKbeta; IKK[b]
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>MR210500 representing NM_001159774
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGCATCGCC**

ATGAGCTGGTCACCGTCCCTCCCAACCCAGACATGTGGAGCCTGGGAAATGAAAGAACGCCTGGGGACCG
 GGGGATTTGGAAACGTCAATCCGGTGGCACAATCAGGCGACAGGTGAACAGATCGCCATCAAGCAATGCCG
 ACAGGAGCTCAGCCCAAAGAACAGAGACCGCTGGTGCCTCGAAATCCAGATCATGAGAAGGCTGAACCAT
 CCCAATGTGGTGGCTGCCCGGGATGTCCAGAGGGGATGCAGAACCTGGCACCCAATGATTTGCCACTGC
 TGGCCATGGAGTACTGCCAAGGAGGAGATCTCCGAAGATACTTGAACCAGTTCGAGAAGTCTGTGGCCT
 GCGGGAAGGAGCTGTCTTACCCTGCTGAGTGACATCGCATCGGCTCTTAGATACCTTACGAAAACAGA
 ATCATCCATCGAGACCTGAAGCCAGAAAACATCGTTCTGCAGCAAGGAGAGAAAAGATTAATACACAAAA
 TTATTGATCTAGGATATGCCAAGGAGCTGGATCAGGGCAGTCTGTGCACGTCATTTGTGGGACTCTGCA
 ATACCTGGCGCCAGAGCTTCTGGAGCAGCAGAAGTACACCGTGACCGTTGACTACTGGAGCTTCGGCACC
 CTGGCCTTCGAGTGCATCACTGGCTTCCGGCCCTTCTCCCTAACTGGCAGCCTGTGCAGTGGCACTCCA
 AAGTCCGGCAGAAGAGCGAAGTGGACATCGTTGTTAGTGAAGACTTGAATGGAGCAGTGAAGTTTCAAG
 TTCGCTACCCTTCCCAATAATCTTAACAGTGTCTTGGCTGAACGGCTGGAGAAGTGGTGCAGCTGATG
 CTTATGTGGCACCTCGGCAAAGGGGCACGGATCCCCAGTATGGCCCCAACGGCTGCTTCAGAGCCCTGG
 ATGACATCTTGAAGCTGGTTCATGTCTTGAACATGGTTCACAGGCACCGTTCACACATACCCCGT
 GACGGAGGATGAGAGTCTGCAGAGCTTAAAAACCAGAATCCAGGAAGACACGGGGATCCTGGAGACAGAC
 CAGGAGCTGCTGCAAGAGGCAGGGCTGGTGTCTCCCTGACAAGCCTGCTACTCAGTGCATCTCAGACA
 GCAAGACAAACGAGGGCCCTCACGTTGGACATGGATCTTGTCTTTTCTCTTTGACAACAGTAAAATCAACT
 TGAGACTCAGATCACCCCGACCCCAACCGGAAAGTGTAGCTGTATCCTTCAGGAGCCCAAGCGGAAC
 CTCTCCTTCTTCCAGCTGAGGAAAGTGTGGGGCCAAGTCTGGCACAGCATCCAGACGCTGAAGGAAGACT
 GTAACCGGCTGCAGCAGGGACAGCGAGCAGCCATGATGAGTCTCCTCCGGAATAACAGCTGCCTCTCTAA
 GATGAAGAACGCCATGGCTCCACGGCCAGCAGCTCAAGGCCAAGCTGGACTTCTTCAAAACCAGCATC
 CAGATCGACCTGGAGAAGTATAAAGAGCAGACCGAGTTTGGGATCACCTCAGATAAATTGCTGCTGGCTT
 GCGGGGAGATGGAGCAGGCTGTGGAGCAGTGTGGCGGGAGAATGACGTGAAGCATCTAGTAGAGCGGAT
 GATGGCACTGCAGACTGACATTGTGGACCTGCAGAGGAGCCGATGGGTGGAAGCAGGGGGGCACCCTG
 GATGACCTAGAGGAACAAGCGAGGGAGCTCTACCGAAGACTCAGGAGAAAGCAAGAGACCAAGGACAG
 AAGGTGACAGCCAGGAGATGGTACGGCTGCTGCTCAGGCAATCCAAAGCTTTGAGAAGAAAGTTGGGT
 GATTTATACACAGCTCAGTAAGACCGTGGTTTGTAAAGCAGAAGGCACTGGAGTTGCTGCCAAGGTAGAA
 GAGGTAGTGAGCCTTATGAACGAGGACGAGAGGACCGTGGTCCGGCTTCAGGAGAAGCGGCAAGGAAC
 TCTGGAACCTCCTGAAGATCGCTGTAGCAAAGTCCGAGGTCCCGTGAAGTGAAGCCAGACAGCATGAA
 TGTGTCTCGACTCAGTACCCTGGTCACTAATGTCCCAGCCTTCCAGTGCCTGTGACAGCTTACCTGAA
 TCAGACAAGAAAAGTGAAGAAGTGGTGGCCGAAGCCACGCCCTGCTCCCGGCTAGAAAAGTGGCTGC
 AGGACTGTGAAGGAGCAAGACAGAAGCTTACGACTCTAGACTGGAGCTGGTTACAGATGGAGGATGA
 AGAAAGGTGTAGCCTGGAGCAGGCTGTGAC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR210500 representing NM_001159774
 Red=Cloning site Green=Tags(s)


MSWSPSLPTQTCGAWEMKERLGTGGFGNVIRWHNQATGEQIAIKQCRQELSPKNRDRWCLEIQIMRRLNH
 PNVVAARDVPEGMQNLAPNDLPLLAMEYCQGGDLRRYLNQFENCCLREGAVLTLLSDIASALRYLHENR
 I IHRDLKPENIVLQQGEKRL IHKIIDLGYAKELDQGLCTSFVGTLQYLAPELLEQQKYTVVDYWSFGT
 LAFECITGFRPFLPNWQPVQWHSKVRQKSEVDIVVSEDLNGAVKFSSSLPFPNNLNSVLAERLEKWLQLM
 LMWHPRQRGTDPOYPNGCFRALDDILNLKLVHVLNMVTGTVHTYPTVEDESLQSLKTRIQEDTGILETD
 QELLQEAGLVLLPDKPATQCISDSKTNEGLTLDMDLVFLFDNSKINYETQITPRPQESVSCILQEPKRN
 LSFFQLRKVWGQVWHSIQTLKEDCNRLQQGQRAAMSSLRNNSLSKMKNAMASTAQQLKAKLDFFKTSI
 QIDLEKYEQTEFGITSDKLLLAWREMEQAVEQCQGRENDVKHLVERMMALQTDIVDLQRSPMGRKQGGTL
 DDLEEQARELYRRLREKPRDQRTEGDSQEMVRLLLQAIQSFEEKVRVIYTLQSKTVVCKQKALELLPKVE
 EVVSLMNEDERTVRLQEKQKELWNLLKIACSKVRGPVSGPDSMNVSRLSHPGQLMSQPSSACDSLPE
 SDKKSEELVAEHALCSRLESALQDTVKEQDRSFTTLDWSWLQMEDEERCSLEQACD

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

Sgfl-MluI

Cloning Scheme:



* The last codon before the Stop codon of the ORF

ACCN: NM_001159774

ORF Size: 2271 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001159774.1](#), [NP_001153246.1](#)

RefSeq Size: 3598 bp

RefSeq ORF: 2274 bp


Locus ID: 16150

Cytogenetics: 8 A2

MW: 86.8 kDa

Gene Summary:

Serine kinase that plays an essential role in the NF-kappa-B signaling pathway which is activated by multiple stimuli such as inflammatory cytokines, bacterial or viral products, DNA damages or other cellular stresses. Acts as part of the canonical IKK complex in the conventional pathway of NF-kappa-B activation and phosphorylates inhibitors of NF-kappa-B on 2 critical serine residues. These modifications allow polyubiquitination of the inhibitors and subsequent degradation by the proteasome. In turn, free NF-kappa-B is translocated into the nucleus and activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis. In addition to the NF-kappa-B inhibitors, phosphorylates several other components of the signaling pathway including NEMO/IKBKG, NF-kappa-B subunits RELA and NFKB1, as well as IKK-related kinases TBK1 and IKBKE. IKK-related kinase phosphorylations may prevent the overproduction of inflammatory mediators since they exert a negative regulation on canonical IKKs. Phosphorylates FOXO3, mediating the TNF-dependent inactivation of this pro-apoptotic transcription factor. Also phosphorylates other substrates including NCOA3, BCL10 and IRS1. Within the nucleus, acts as an adapter protein for NFKBIA degradation in UV-induced NF-kappa-B activation. [UniProtKB/Swiss-Prot Function]

Product images:

Circular map for MR210500