

# Product datasheet for MR210501L4V

## Ipo4 (BC003469) Mouse Tagged ORF Clone Lentiviral Particle

## **Product data:**

#### **Product Type: Lentiviral Particles Product Name:** Ipo4 (BC003469) Mouse Tagged ORF Clone Lentiviral Particle Symbol: lpo4 RanBP4, Imp4a Synonyms: **Mammalian Cell** Puromycin Selection: Vector: pLenti-C-mGFP-P2A-Puro (PS100093) mGFP Tag: ACCN: BC003469 ORF Size: 2274 bp The ORF insert of this clone is exactly the same as(MR210501). **ORF** Nucleotide Sequence: **OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info **OTI** Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. **RefSeq:** BC003469.1 **RefSeq Size:** 2833 bp **RefSeq ORF:** 2276 bp Locus ID: 75751 Cytogenetics: 14 28.19 cM



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### CRIGENE Ipo4 (BC003469) Mouse Tagged ORF Clone Lentiviral Particle – MR210501L4V

Gene Summary: Functions in nuclear protein import as nuclear transport receptor. Serves as receptor for nuclear localization signals (NLS) in cargo substrates. Is thought to mediate docking of the importin/substrate complex to the nuclear pore complex (NPC) through binding to nucleoporin and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to the importin, the importin/substrate complex dissociates and importin is re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran. The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus (By similarity). Mediates the nuclear import of RPS3A. Acts as chaperone for exposed basic domains.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US