

## Product datasheet for MR210661L3

### Dgkg (NM\_138650) Mouse Tagged Lenti ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Dgkg (NM_138650) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Dgkg
Synonyms:	90kDa; 2900055E17Rik; AI854428; Dagk3; E430001K23Rik; mKIAA4131
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR210661).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:




\* The last codon before the Stop codon of the ORF.

ACCN:	NM_138650
ORF Size:	2364 bp



<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>Note:</b>	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
<b>RefSeq:</b>	<a href="#">NM_138650.1</a>
<b>RefSeq Size:</b>	5521 bp
<b>RefSeq ORF:</b>	2367 bp
<b>Locus ID:</b>	110197
<b>UniProt ID:</b>	<a href="#">Q91WG7</a>
<b>Cytogenetics:</b>	16 13.37 cM
<b>Gene Summary:</b>	Diacylglycerol kinase that converts diacylglycerol/DAG into phosphatidic acid/phosphatidate/PA and regulates the respective levels of these two bioactive lipids (PubMed:32033984). Thereby, acts as a central switch between the signaling pathways activated by these second messengers with different cellular targets and opposite effects in numerous biological processes (PubMed:32033984). Has no apparent specificity with regard to the acyl compositions of diacylglycerol (By similarity). Specifically expressed in the cerebellum where it controls the level of diacylglycerol which in turn regulates the activity of protein kinase C gamma (PubMed:32033984). Through protein kinase C gamma, indirectly regulates the dendritic development of Purkinje cells, cerebellar long term depression and ultimately cerebellar motor coordination (PubMed:32033984).[UniProtKB/Swiss-Prot Function]

## Product images:



Circular map for MR210661L3