

Product datasheet for MR210704L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Rasgrp1 (NM_011246) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Rasgrp1 (NM_011246) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Rasgrp1
Synonyms: Rasgrp

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_011246

 ORF Size:
 2385 bp

ORF Nucleotide

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(MR210704).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 011246.2</u>

 RefSeq Size:
 5155 bp

 RefSeq ORF:
 2388 bp

 Locus ID:
 19419

 UniProt ID:
 Q9Z1S3

 Cytogenetics:
 2 59.19 cM







Gene Summary:

Functions as a calcium- and diacylglycerol (DAG)-regulated nucleotide exchange factor specifically activating Ras through the exchange of bound GDP for GTP. Activates the Erk/MAP kinase cascade. Regulates T-cell/B-cell development, homeostasis and differentiation by coupling T-lymphocyte/B-lymphocyte antigen receptors to Ras. Regulates NK cell cytotoxicity and ITAM-dependent cytokine production by activation of Ras-mediated ERK and JNK pathways (By similarity). Functions in mast cell degranulation and cytokine secretion, regulating FcERI-evoked allergic responses (PubMed:17190838). May also function in differentiation of other cell types. Proto-oncogene, which promotes T-cell lymphomagenesis when its expression is deregulated (PubMed:15829980, PubMed:17210708).[UniProtKB/Swiss-Prot Function]