

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for MR210777L4V

## Lrrc8a (BC048152) Mouse Tagged ORF Clone Lentiviral Particle

### **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Lrrc8a (BC048152) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Lrrc8a
Synonyms:	mKIAA1437, MGC61242
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	BC048152
ORF Size:	2430 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR210777).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>BC048152</u>
RefSeq Size:	4344 bp
RefSeq ORF:	2432 bp
Locus ID:	241296
Cytogenetics:	2 B



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** Lrrc8a (BC048152) Mouse Tagged ORF Clone Lentiviral Particle – MR210777L4V

Gene Summary:Essential component of the volume-regulated anion channel (VRAC, also named VSOAC<br/>channel), an anion channel required to maintain a constant cell volume in response to<br/>extracellular or intracellular osmotic changes (PubMed:29769723). The VRAC channel<br/>conducts iodide better than chloride and can also conduct organic osmolytes like taurine (By<br/>similarity). Mediates efflux of amino acids, such as aspartate and glutamate, in response to<br/>osmotic stress (By similarity). Required for channel activity, together with at least one other<br/>family member (LRRC8B, LRRC8C, LRRC8D or LRRC8E); channel characteristics depend on the<br/>precise subunit composition. Can form functional channels by itself (in vitro) (By similarity).<br/>Involved in B-cell development: required for the pro-B cell to pre-B cell transition<br/>(PubMed:14660746, PubMed:24752297). Also required for T-cell development<br/>(PubMed:24752297).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US