

## Product datasheet for MR210799L4

### Cd177 (NM\_026862) Mouse Tagged Lenti ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Cd177 (NM_026862) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Cd177
Synonyms:	1190003K14Rik; Pdp3
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR210799).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


ACCN:	NM_026862
ORF Size:	2451 bp



[View online »](#)

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>Note:</b>	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
<b>RefSeq:</b>	<a href="#">NM_026862.3</a> , <a href="#">NP_081138.2</a>
<b>RefSeq Size:</b>	2733 bp
<b>RefSeq ORF:</b>	2454 bp
<b>Locus ID:</b>	68891
<b>UniProt ID:</b>	<a href="#">Q8R2S8</a>
<b>Cytogenetics:</b>	7 A3
<b>Gene Summary:</b>	In association with beta-2 integrin heterodimer ITGAM/CD11b and ITGB2/CD18, mediates activation of TNF-alpha primed neutrophils including degranulation and superoxide production (By similarity). In addition, by preventing beta-2 integrin internalization and attenuating chemokine signaling favors adhesion over migration (By similarity). Heterophilic interaction with PECAM1 on endothelial cells plays a role in neutrophil transendothelial migration in vitro (By similarity). However, appears to be dispensable for neutrophil recruitment caused by bacterial infection in vivo (PubMed:25359465). Acts as a receptor for the mature form of protease PRTN3 allowing its display at the cell surface of neutrophils (By similarity). By displaying PRTN3 at the neutrophil cell surface, may play a role in enhancing endothelial cell junctional integrity and thus vascular integrity during neutrophil diapedesis (By similarity).[UniProtKB/Swiss-Prot Function]

## Product images:



Circular map for MR210799L4