

## Product datasheet for MR210905L4

### Lztr1 (NM\_025808) Mouse Tagged Lenti ORF Clone

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Expression Plasmids  |
| Product Name:             | Lztr1 (NM_025808) Mouse Tagged Lenti ORF Clone                 |
| Tag:                      | mGFP   |
| Symbol:                   | Lztr1  |
| Synonyms:                 | 1200003E21Rik; AI591627; AW550890; TCFL                        |
| Mammalian Cell Selection: | Puromycin  |
| Vector:                   | pLenti-C-mGFP-P2A-Puro (PS100093)                              |
| E. coli Selection:        | Chloramphenicol (34 ug/mL)                                     |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(MR210905). |
| Restriction Sites:        | SgfI-MluI  |
| Cloning Scheme:           |  |

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


|           |           |
|-----------|-----------|
| ACCN:     | NM_025808 |
| ORF Size: | 2511 bp   |



[View online »](#)

|                               |   |
|-------------------------------|---|
| <b>OTI Disclaimer:</b>        | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>  |
| <b>OTI Annotation:</b>        | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.  |
| <b>Components:</b>            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).  |
| <b>Reconstitution Method:</b> | <ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol> |
| <b>RefSeq:</b>                | <a href="#">NM_025808.2</a> , <a href="#">NP_080084.2</a>   |
| <b>RefSeq Size:</b>           | 3108 bp   |
| <b>RefSeq ORF:</b>            | 2514 bp   |
| <b>Locus ID:</b>              | 66863   |
| <b>UniProt ID:</b>            | <a href="#">Q9CQ33</a>  |
| <b>Cytogenetics:</b>          | 16 A3   |
| <b>Gene Summary:</b>          | This gene encodes a member of the BR-C, ttk and bab-kelch superfamily that, in humans, localizes to the Golgi network and is associated with the ras / mitogen-activated protein kinase pathway. Loss-of-function mutations in the human ortholog are associated with glioblastoma multiforme, schwannomatosis, Noonan syndrome, and DiGeorge syndrome. [provided by RefSeq, Sep 2016]  |

Product images:



Circular map for MR210905L4