

## Product datasheet for **MR211034L3V**

### Copg1 (NM\_017477) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Lentiviral Particles   |
| Product Name:             | Copg1 (NM_017477) Mouse Tagged ORF Clone Lentiviral Particle   |
| Symbol:                   | Copg1  |
| Synonyms:                 | AU019265; BC056168; Copg; D6Erttd71e; D6Wsu16e   |
| Mammalian Cell Selection: | Puromycin  |
| Vector:                   | pLenti-C-Myc-DDK-P2A-Puro (PS100092)   |
| Tag:                      | Myc-DDK  |
| ACCN:                     | NM_017477  |
| ORF Size:                 | 2625 bp  |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(MR211034).   |
| OTI Disclaimer:           | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a> |
| OTI Annotation:           | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| RefSeq:                   | <a href="#">NM_017477.2</a> , <a href="#">NP_059505.1</a>  |
| RefSeq Size:              | 4014 bp  |
| RefSeq ORF:               | 2625 bp  |
| Locus ID:                 | 54161  |
| UniProt ID:               | <a href="#">Q9QZE5</a>   |
| Cytogenetics:             | 6 39.13 cM   |



[View online »](#)

**Gene Summary:**

The coatomer is a cytosolic protein complex that binds to dilysine motifs and reversibly associates with Golgi non-clathrin-coated vesicles, which further mediate biosynthetic protein transport from the ER, via the Golgi up to the trans Golgi network. Coatomer complex is required for budding from Golgi membranes, and is essential for the retrograde Golgi-to-ER transport of dilysine-tagged proteins. In mammals, the coatomer can only be recruited by membranes associated to ADP-ribosylation factors (ARFs), which are small GTP-binding proteins; the complex also influences the Golgi structural integrity, as well as the processing, activity, and endocytic recycling of LDL receptors (By similarity). Required for limiting lipid storage in lipid droplets. Involved in lipid homeostasis by regulating the presence of perilipin family members PLIN2 and PLIN3 at the lipid droplet surface and promoting the association of adipocyte triglyceride lipase (PNPLA2) with the lipid droplet surface to mediate lipolysis. [UniProtKB/Swiss-Prot Function]