

# **Product datasheet for MR211131L3V**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Hip1 (NM\_146001) Mouse Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type: Lentiviral Particles

**Product Name:** Hip1 (NM\_146001) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Hip<sup>2</sup>

**Synonyms:** 2610109B09Rik; A930014B11Rik; E130315I21Rik; mKIAA4113

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_146001

 ORF Size:
 3087 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(MR211131).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 146001.2</u>, <u>NP 666113.2</u>

 RefSeq Size:
 7883 bp

 RefSeq ORF:
 3090 bp

 Locus ID:
 215114

 UniProt ID:
 Q8VD75

 Cytogenetics:
 5 75.18 cM







#### **Gene Summary:**

Plays a role in clathrin-mediated endocytosis and trafficking (PubMed:11577110). Involved in regulating AMPA receptor trafficking in the central nervous system in an NMDA-dependent manner (PubMed:12839988, PubMed:17329427). Regulates presynaptic nerve terminal activity (PubMed:17928447). Enhances androgen receptor (AR)-mediated transcription (By similarity). May act as a proapoptotic protein that induces cell death by acting through the intrinsic apoptosis pathway (By similarity). Binds 3-phosphoinositides (via ENTH domain) (By similarity). May act through the ENTH domain to promote cell survival by stabilizing receptor tyrosine kinases following ligand-induced endocytosis (By similarity). May play a functional role in the cell filament networks (By similarity). May be required for differentiation, proliferation, and/or survival of somatic and germline progenitors (PubMed:11604514, PubMed:14998932, PubMed:16967501, PubMed:17928447).[UniProtKB/Swiss-Prot Function]