

Product datasheet for MR211142L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Gria1 (NM_008165) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Gria1 (NM_008165) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Gria1

Synonyms: 2900051M01Rik; Glr-1; GluA1; Glur-1; GluR-A; gluR-K1; Glur-1; GluRA; HIPA1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_008165 **ORF Size:** 2724 bp

ORF Nucleotide

_. _.

Sequence:

The ORF insert of this clone is exactly the same as(MR211142).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 008165.2

 RefSeq Size:
 5755 bp

 RefSeq ORF:
 2724 bp

 Locus ID:
 14799

Cytogenetics: 11 34.51 cM







Gene Summary:

lonotropic glutamate receptor. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. In the presence of CACNG4 or CACNG7 or CACNG8, shows resensitization which is characterized by a delayed accumulation of current flux upon continued application of glutamate.[UniProtKB/Swiss-Prot Function]